

LONGEVITY OF RESEEDED GRASS SPECIES USED FOR RESTORING THE DEGRADED SUBALPINE MEADOWS

Teodor MARUȘCA^{1,2}, Vasile A. BLAJ¹, Vasile MOCANU¹

Abstract. *In the summer of 1996, a degraded grassland, invaded by *Nardus stricta* species, located at 1800 m altitude from subalpine level of the Bucegi Mountains, after total herbicide with glyphosate, liming using CaO at 2/3 Ah (in the autumn of 1995) and paddocking with sheep (5 nights, 1 sheep / m²) has been over-seeded or reseeded. The grass seed mixture was composed of *Phleum pratense* 40%; *Festuca pratensis* 25%; *Lolium perenne* 5%, *Lotus corniculatus* 15% and *Trifolium hybridum* 15%. A part of the variants have been fertilized with chemical fertilizers with doses of N 150 K P50 50 kg / ha and others plots have been fertilized with organic fertilizer by a paddocking system applied before of reseeded grassland establishment. In 2004 and 2011 an organic fertilizing by cattle paddocking, has been practiced. The reseeded species that do not reach maturity remain a much longer period of time than is known in the technical literature, this being 2-3 times higher in the high mountains than in the lowlands and hills. In the grassy carpet the *Phleum pratense* species survives in large proportion, even after 20 years of sowing.*

Keywords: degraded grassland, fertilizer, herbicide, over-seeding, reseeded.

INTRODUCTION

The mountain pastures, used by grazing for many years, are exposed to processes of degradation of floristic composition, if not properly maintained, fertilized and used rationally [1].

By applying the organic (paddocking system) and chemical fertilizers, degraded mountain grasslands, invaded by strict *Nardus stricta* species, can turn into valuable pastures, dominated by *Festuca rubra* and *Agrostis capillaris* [4]

¹ Research and Development Institute for Grasslands, 5 Cucului Str., 500128, 0268472701; fax 0268475295, www.pajisti-grassland.ro, office@pajisti-grassland.ro

² Academy of Romania Scientists, Splaiul Independenței 54, 050094, Bucharest, Romania

One of the most effective ways to improve *Nardus stricta* degraded grasslands is by total sward destruction (herbicides, harrowing, milling, plowing, etc.), followed by the establishment of reseeded pastures with high quality and production [2]

Such research work to improve more effectively the *Nardus stricta* grasslands has been conducted in sub-alpine altitudinal level in the last 20 years [3].

One such long-term research activity in sub-alpine level was possible to determine the survival duration of some species of perennial sown grasses that it is different from the conditions at lower altitudinal levels, thematic presented in this paper.

MATERIAL AND METHOD

At the Research Mountain Grasslands Base from the Bucegi Massive, located at 1,800 m of altitude on a degraded pasture, with participation of *Nardus stricta* species over 60%, in the autumn of 1996, the grassy carpet was sprayer with 5 L/ha Roundup product (glyphosate). After two weeks were applied 7.5 tons per ha of lime (CaO) to correct 2/3 Ah of soil acidity (hydrolytic acidity).

In June 1996, after sheep climbing on the mountain, it paddocked a sheep 5 nights/1 m², after which the land was harrowing at 2-3 cm deep and sown with a mixture of: *Phleum pratense* 40% *Festuca pratensis* 25% *Lolium perenne* 5%, *Lotus corniculatus* 15% and *Trifolium hybridum* 15%, all missing species of wild flora.

In the organic fertilized variant, paddocking was repeated, at the same rate, with sheep in 2004, and with cows in 2012. Similarly, for comparison, the same mixture of grasses with chemical fertilizers N 150 P 50 K 50 kg / ha, during 1996-1998, 2004-2006 and 2012-2014, was fertilized.

After sampling for dry matter (DM) and chemical analysis, pasture was used by grazing with cows. The sown species were not able to multiply by self-seeding, because they have not reached maturity. Annually, between 1997-2016, flora observations using the KLAPP-ELEMBERG method of participation percentage in phytomass, were made.

RESULTS AND DISCUSSION

Following the floristic observations, over 20 years, every two years, it was possible an overview of the dynamics of the sown and spontaneous species in the chemical fertilized variant (Table 1.).

From this data follows the best representation of sown species of 86% was on 1999-2000 and the lowest of 21% in 2011-2012. The basic *Phleum pratense* species was maintained at a rate of 21-66% in grassy carpet, being the only species that has resisted for 20 years. The remaining species have disappeared at

different times, namely: *Festuca pratensis* after 12 years, *Lolium perenne* after 4 years, *Trifolium hybridum* after 10 years and *Lotus corniculatus* after 8 years.

In the grassy carpet the *Poa pratensis* species appears, from spontaneous flora, only after 15 years from the start of the improvement work and *Taraxacum officinale* after 7 years.

The pastoral value of reseeded pastures, chemically fertilized, is 61-83 and it is good to very good. On variant organic fertilized by paddocking system, the success of sown species is better than chemical fertilization, being 90% in the period 1997-1998 and 22% in 2015-2016 (Table 2).

In the 20 years of observations, the only species that survived in this case was the *Phleum pratense* species, in a variable proportion of 22-55%. In this case *Festuca pratensis* disappeared after 16 years, *Lolium perenne* after 6 years and the *Trifolium hybridum* species after 10 years, a little more enduring than the chemically fertilized variant.

Instead, in the paddocked variant, in the first year, from spontaneous flora, the *Taraxacum officinale* has established and after the 7th year *Poa pratensis* has maintained around 5%. The pastoral value of the organic variant has indices between 74 and 90, being superior to the chemical alternative.

A comparison of the influence of the type of chemical to organic fertilizer is presented in Table 3. These data show that the organic fertilization has favored the sown species *Phleum pratense*, *Festuca pratensis* and *Trifolium hybridum* and the spontaneous species *Trifolium repens* and *Poa pratensis*. The viability of the sown species, fertilized organically is longer than the chemical variant, by 2 years for *Lolium perenne* and 4 years for *Festuca pratensis*.

Similarly, in the organic variant, the *Poa pratensis* species appears faster by 8 years and *Taraxacum officinale* by 6 years compared to the chemical solution. In general, the index of forage value of organic variant is higher by 9 than the chemical variant.

Table 1
The evolution of the floristic composition of sown pastures, fertilized chemically, Blana- Bucegi Mountains 1997 - 2016
(% Participation in aboveground phytomass)

Specii	1997- 1998	1999- 2000	2001- 2002	2003- 2004	2005- 2006	2007- 2008	2009- 2010	2011- 2012	2013- 2014	2015- 2016
SOWN SPECIES	64	86	53	57	55	37	29	21	34	24
Perennial grasses										
<i>Phleum pratense</i>	30	66	41	45	49	33	29	21	34	24
<i>Festuca pratensis</i>	12	4	2	2	4	4
<i>Lolium perenne</i>	9	+
Perennial forage legumes										
<i>Trifolium hybridum</i>	12	16	10	10	2
<i>Lotus corniculatus</i>	1	+	+	+
SPONTANEOUS SPECIES	36	14	47	43	45	63	71	79	66	76
Perennial grasses										
<i>Agrostis capillaris</i>	8	4	15	14	17	33	34	27	20	23
<i>Festuca nigrescens</i>	7	1	12	8	5	6	8	11	14	9
<i>Poa pratensis</i>	1	3	4
Alte graminee	16	3	7	10	9	11	15	21	13	21
Perennial forage legumes										
<i>Trifolium repens</i>	4	4	10	6	7	10	11	10	7	10
Others families										
<i>Ligusticum mutelina</i>	+	1	+	+	1	2	2	5	3	4
<i>Potentilla ternata</i>	1	+	+	1	1	1	1	2	5	2
<i>Taraxacum officinale</i>	.	.	.	+	2	+	+	1	+	+
Alte specii	+	1	3	4	3	+	+	1	1	3
PASTORAL VALUE	77	83	77	76	78	73	68	61	67	62

Table 2
 The evolution of the floristic composition of sown pastures, organic fertilized by paddocking, Blana- Bucegi Mountains, 1997 - 2016
 (% Participation in aboveground phytomass)

Species	1997- 1998	1999- 2000	2001- 2002	2003- 2004	2005- 2006	2007- 2008	2009- 2010	2011- 2012	2013- 2014	2015- 2016
SOWN SPECIES	90	76	52	56	67	53	29	24	24	22
Perennial grasses										
<i>Phleum pratense</i>	22	33	35	45	55	42	25	23	24	22
<i>Festuca pratensis</i>	20	7	3	6	10	11	4	1	.	.
<i>Lolium perenne</i>	9	1	+
Perennial forage legumes										
<i>Trifolium hybridum</i>	38	35	14	5	2
<i>Lotus corniculatus</i>	1	+	+
SPONTANEOUS SPECIES	10	24	48	44	33	47	71	76	76	78
Perennial grasses										
<i>Agrostis capillaris</i>	1	8	12	13	17	28	34	32	20	25
<i>Festuca nigrescens</i>	1	1	1	5	1	+	1	1	2	5
<i>Poa pratensis</i>	.	.	.	2	3	4	5	4	5	5
Other grasses	2	1	14	8	2	1	5	8	5	8
Perennial forage legumes										
<i>Trifolium repens</i>	6	13	14	12	8	8	16	22	30	24
Others families										
<i>Taraxacum officinale</i>	+	+	1	2	1	2	5	4	7	5
<i>Ligusticum mutelina</i>	+	1	2	1	1	2	3	3	3	3
<i>Potentilla ternata</i>	+	+	1	1	+	1	1	1	2	1
Others species	+	+	3	+	+	1	1	1	2	1
PASTORAL VALUE	90	88	76	82	89	84	76	75	79	74

Table 3
Comparative data on the effect of fertilizer on species type and their longevity in the grassy carpet Blana - Bucegi 1997 – 2016

Specii	Semănat 1996	Media 1997 - 2016		Dif. + -		Durată specie (ani)		
		Chimic	Organic	Chimic	Organic	Chimic	Organic	Dif. + -
SOWN SPECIES	100	46,0	49,3	+3,3				
Perennial grasses								
<i>Phleum pratense</i>	40	37,2	32,6	-4,6		20	20	0
<i>Festuca pratensis</i>	25	2,8	6,2	+3,4		12	16	+4
<i>Lolium perenne</i>	5	0,9	1,0	+0,1		4	6	+2
Perennial forage legumes								
<i>Trifolium hybridum</i>	15	5,0	9,4	+4,4		10	10	0
<i>Lotus corniculatus</i>	15	0,1	0,1	0		8	6	-2
SPONTANEOUS SPECIES	0	54,0	50,7	-3,3				
Perennial grasses								
<i>Agrostis capillaris</i>	X	19,5	19,0	-0,5		20	20	0
<i>Festuca nigrescens</i>	X	8,1	1,9	-6,2		20	20	0
<i>Poa pratensis</i>	X	0,8	2,8	+2,0		6	14	+8
Other grasses	X	12,6	5,4	-7,2		20	20	0
Perennial forage legumes								
<i>Trifolium repens</i>	X	7,9	15,3	+7,4		20	20	0
Others families								
<i>Ligusticum mutelina</i>	X	1,8	1,9	+0,1		20	20	0
<i>Potentilla ternata</i>	X	1,4	0,8	-0,6		20	20	0
<i>Taraxacum officinale</i>	X	0,3	2,7	+2,4		14	20	+6
Others species	X	1,6	0,9	-0,7		20	20	0
PASTORAL VALUE	94	72	81	+9		X	X	X

CONCLUSIONS

- (1) In the sub-alpine high mountains are few species of perennial grasses and forage legumes adapted to these soil and climate conditions, less favorable to plant growth;
- (2) By providing optimal conditions of soil and fertilizer, timothy (*Phleum pratense*) manages to survive more than 20 years at a rate of 37% in the chemical fertilizer variant and 33% in organic fertilization (paddock) compared to 40% at sowing;
- (3) The longevity of perennial sown grasses and legumes at high mountain altitude is at least double than those sown at lower altitude in the plains;
- (4) Paddock by animals has favored the sown species *Festuca pratensis* and *Trifolium hybridum* compared to the chemical fertilization that has favored the *Phleum pratense* species;
- (5) From spontaneous flora *Poa pratensis*, *Trifolium repens* and *Taraxacum officinale* are stimulated by paddocking and *Festuca nigrescens* by chemical fertilization.

REFERENCES

- [1] Bărbulescu, Motcă Gh., 1983, *Pășunile munților înalți*, Ed. Ceres, București
- [2] Marușca T., 1977, *Sisteme de înființare a pajiștilor temporare pe suprafețele dominate de *Nardus stricta**, Lucrări științifice ale SCCP Măgurele – Brașov, vol. III, pp. 35-49, Redacția material de propagandă agricolă, București
- [3] Marușca T., Andreoiu Andreea, Blaj V.A., Mocanu V., Haș C.E, *Long term effect of improvement methods on subalpine degraded *Nardus stricta* l. grasslands*, 2015, Academy of Romanian Scientists, Annals Series on Agriculture, Silviculture and Veterinary Medicine Sciences, vol. 4. nr 1-2, București, pp.42-49, ISSN 2069 - 1149
- [4] Pușcaru D., Pușcaru-Soroceanu E., Paucă A., Șerbănescu I., Beldie Al., Ștefureac Tr., Cernescu N., Saghin N., Crețu V., Lupan L., Tașcenco V., 1956, *Pășunile alpine din Munții Bucegi*, Ed. Academiei Republicii Populare Române