

# FREDHOLM TOEPLITZ OPERATORS ON THE WEIGHTED BERGMAN SPACES\*

Namita Das<sup>†</sup> Swarupa Roy<sup>‡</sup>

DOI <https://doi.org/10.56082/annalsarscimath.2021.1-2.178>

## Abstract

In this paper we have shown that if  $\phi \in (L_h^2(dA_\alpha))^\perp \cap L^\infty(\mathbb{D})$  and  $\text{Range}T_\phi^{(\alpha)}$  is closed, then the Toeplitz operator  $T_\phi^{(\alpha)} \in \mathcal{L}(L_a^2(dA_\alpha))$  is a Fredholm operator of index zero and  $T_\phi^{(\alpha)}$  is not of finite rank. Several applications of the result were also obtained. We further show that if  $\phi \in L_{M_n}^\infty(\mathbb{D})$  is such that  $T_\phi$  is Fredholm and of index zero in  $\mathcal{L}(L_a^{2,\mathbb{C}^n}(dA_\alpha))$  then there exists  $\psi \in E_{n \times n} = E \otimes M_n$  such that  $T_{\phi+\delta\psi}$  is invertible for all sufficiently small nonzero  $\delta$ . Here  $E$  is a total subspace of  $L^\infty(\mathbb{D})$  and  $M_n$  is the set of all  $n \times n$  matrices with complex entries.

MSC: 47B38, 47B32

**keywords:** Weighted Bergman spaces, Finite rank operator, Toeplitz operator, Little Hankel operator, Bounded harmonic functions.

## 1 Introduction

Let  $dA(z) = \frac{1}{\pi} dx dy = \frac{1}{\pi} r dr d\theta$  be the normalized area measure on the open unit disk  $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$  in the complex plane  $\mathbb{C}$ . For  $\alpha > -1$ , let

---

\*Accepted for publication on March 30-th, 2021

<sup>†</sup>[namitadas440@yahoo.co.in](mailto:namitadas440@yahoo.co.in) P. G. Dept. of Mathematics, P. G. Dept. of Mathematics, Utkal University, Vani Vihar, Utkal University, Vani Vihar, Bhubaneswar- 751004, Odisha, India

<sup>‡</sup>[swarupa.roy@gmail.com](mailto:swarupa.roy@gmail.com) P. G. Dept. of Mathematics, P. G. Dept. of Mathematics, Utkal University, Vani Vihar, Utkal University, Vani Vihar, Bhubaneswar- 751004, Odisha, India

$dA_\alpha(z) = (\alpha + 1)(1 - |z|^2)^\alpha dA(z)$ ,  $z \in \mathbb{D}$ . Let  $L^2(\mathbb{D}, dA_\alpha)$  be the space of all absolutely square-integrable Lebesgue measurable functions on  $\mathbb{D}$ . Let the weighted Bergman space of the disk  $\mathbb{D}$ ,  $L_a^2(dA_\alpha)$  be the subspace consisting of all analytic functions of  $L^2(\mathbb{D}, dA_\alpha)$ . The space  $L^2(\mathbb{D}, dA_\alpha)$  is a Hilbert space with respect to the inner product defined by

$$\langle f, g \rangle = \int_{\mathbb{D}} f(z) \overline{g(z)} dA_\alpha(z), \quad f, g \in L^2(\mathbb{D}, dA_\alpha).$$

The weighted Bergman space  $L_a^2(dA_\alpha)$  is a closed subspace of  $L^2(\mathbb{D}, dA_\alpha)$  [4] and hence a reproducing kernel Hilbert space with the reproducing kernel given by  $K^{(\alpha)}(z, \bar{w}) = \frac{1}{(1 - z\bar{w})^{\alpha+2}}$ ,  $z, w \in \mathbb{D}$ . The orthogonal projection  $P_\alpha$  from the Hilbert space  $L^2(\mathbb{D}, dA_\alpha)$  onto the closed subspace  $L_a^2(dA_\alpha)$  is given by  $P_\alpha f(z) = \int_{\mathbb{D}} K^{(\alpha)}(z, \bar{w}) f(w) dA_\alpha(w)$ . Let  $k_z(w) = \frac{1 - |z|^2}{(1 - w\bar{z})^2}$ . The functions  $k_z^{1+\frac{\alpha}{2}} = \frac{K^{(\alpha)}(z, w)}{\sqrt{K^{(\alpha)}(w, \bar{w})}} = \frac{(1 - |w|^2)^{1+\frac{\alpha}{2}}}{(1 - \bar{w}z)^{2+\alpha}}$  are the normalized reproducing kernels of  $L_a^2(dA_\alpha)$ . The sequence of functions  $\{\frac{z^n}{\gamma_{n,\alpha}}\}$  form an orthonormal basis [13] for  $L_a^2(dA_\alpha)$  where  $\gamma_{n,\alpha}^2 = \|z^n\|^2 = (\alpha + 1) \int_{\mathbb{D}} |z|^{2n} (1 - |z|^2)^\alpha dA(z) = \frac{\Gamma(n+1)\Gamma(\alpha+1)}{\Gamma(n+\alpha+2)} \sim (n+1)^{-\alpha-1}$ . Let  $L^\infty(\mathbb{D}, dA)$  be the space of all essentially bounded Lebesgue measurable functions on  $\mathbb{D}$ . For  $\varphi \in L^\infty(\mathbb{D})$ , define  $\|\varphi\|_\infty = \text{ess sup} \{|\varphi(z)| : z \in \mathbb{D}\} < \infty$ . The space  $L^\infty(\mathbb{D})$  is a Banach space with respect to the  $\|\cdot\|_\infty$ . Let  $H^\infty(\mathbb{D})$  be the space of all bounded analytic functions on  $\mathbb{D}$  and  $h^\infty(\mathbb{D})$  be the space of all bounded harmonic functions on  $\mathbb{D}$ .  $L_h^2(dA_\alpha) = L_a^2(dA_\alpha) \oplus \left(\overline{L_a^2(dA_\alpha)}\right)_0$  where  $\left(\overline{L_a^2(dA_\alpha)}\right)_0 = \{f \in L_a^2(dA_\alpha) : f(0) = 0\}$ .

Given a function  $\varphi \in L^\infty(\mathbb{D})$ , we define an operator  $T_\varphi^{(\alpha)}$  on  $L_a^2(dA_\alpha)$  by  $T_\varphi^{(\alpha)} f = P_\alpha(\varphi f)$ ,  $f \in L_a^2(dA_\alpha)$ . The operator  $T_\varphi^{(\alpha)}$  is called the Toeplitz operator on the weighted Bergman space with symbol  $\varphi$ . Since the projection  $P_\alpha$  has norm 1, we have  $\|T_\varphi^{(\alpha)}\| \leq \|\varphi\|_\infty$ . We can write  $T_\varphi^{(\alpha)}$  as an integral operator as,  $T_\varphi^{(\alpha)} f(z) = \int_{\mathbb{D}} \varphi(w) K^{(\alpha)}(z, w) f(w) dA_\alpha(w) = \int_{\mathbb{D}} \frac{\varphi(w) f(w)}{(1 - z\bar{w})^{\alpha+2}} dA_\alpha(w)$ . Let  $\overline{L_a^2(dA_\alpha)} = \{\bar{f} : f \in L_a^2(dA_\alpha)\}$ . The space  $\overline{L_a^2(dA_\alpha)}$  is a closed subspace of  $L^2(\mathbb{D}, dA_\alpha)$ . The little Hankel operator  $h_\phi^{(\alpha)}$  with symbol  $\phi$  is defined by  $h_\phi^{(\alpha)} f = \overline{P_\alpha(\phi f)}$ ,  $f \in L_a^2(dA_\alpha)$  where  $\overline{P_\alpha}$  is the orthogonal projection from the Hilbert space  $L^2(\mathbb{D}, dA_\alpha)$  onto  $\overline{L_a^2(dA_\alpha)}$ . Clearly,  $\|h_\phi^{(\alpha)}\| \leq \|\phi\|_\infty$ . Define  $J_\alpha$  from  $L^2(\mathbb{D}, dA_\alpha)$  into itself by  $(J_\alpha f)(z) = f(\bar{z})$ ,  $z \in \mathbb{D}$ . The operator  $J_\alpha$  is an unitary operator. For  $\phi \in L^\infty(\mathbb{D})$ ,

define  $S_\phi^{(\alpha)}$  from  $L_a^2(dA_\alpha)$  into itself by  $S_\phi^{(\alpha)}f = P_\alpha(J_\alpha(\phi f))$ . The operator  $S_\phi^{(\alpha)}$  is a linear operator and  $\|S_\phi^{(\alpha)}\| \leq \|\phi\|_\infty$ . It is not difficult to verify that  $h_\phi^{(\alpha)} = J_\alpha S_\phi^{(\alpha)}$ . Thus we shall refer in the sequel, both the operators  $h_\phi^{(\alpha)}$  and  $S_\phi^{(\alpha)}$  as little Hankel operators on  $L_a^2(dA_\alpha)$ .

Let  $L_a^{2,\mathbb{C}^n}(dA_\alpha) = L_a^2(dA_\alpha) \otimes \mathbb{C}^n$  and  $L_{M_n}^\infty(\mathbb{D}) = L^\infty(\mathbb{D}) \otimes M_n$  where  $M_n(\mathbb{C}) = M_n$ ,  $n \geq 1$  is the set of all  $n \times n$  matrices with entries in  $\mathbb{C}$ . The space  $L_a^{2,\mathbb{C}^n}(dA_\alpha)$ ,  $n \geq 1$  is called the vector-valued weighted Bergman space. The inner product on  $L_a^{2,\mathbb{C}^n}(dA_\alpha)$  is defined as

$$\langle f, g \rangle_{L_a^{2,\mathbb{C}^n}(dA_\alpha)} = \int_{\mathbb{D}} \langle f(z), g(z) \rangle_{\mathbb{C}^n} dA_\alpha(z).$$

With this inner product  $L_a^{2,\mathbb{C}^n}(dA_\alpha)$  is a Hilbert space. The norm defined on  $L_a^{2,\mathbb{C}^n}(dA_\alpha)$  is given by

$$\|f\|_{L_a^{2,\mathbb{C}^n}(\mathbb{D}, dA_\alpha)}^2 = \int_{\mathbb{D}} \|f(z)\|_{\mathbb{C}^n}^2 dA_\alpha(z).$$

It is a closed subspace of  $L^{2,\mathbb{C}^n}(\mathbb{D}, dA_\alpha) = L^2(\mathbb{D}, dA_\alpha) \otimes \mathbb{C}^n$ . Let  $P_\alpha^{\mathbb{C}^n}$  denote the orthogonal projection from  $L^{2,\mathbb{C}^n}(\mathbb{D}, dA_\alpha)$  onto  $L_a^{2,\mathbb{C}^n}(dA_\alpha)$ . For  $\phi \in L_{M_n}^\infty(\mathbb{D})$ , we define the Toeplitz operator  $T_\phi$  from  $L_a^{2,\mathbb{C}^n}(dA_\alpha)$  into itself as  $T_\phi f = P_\alpha^{\mathbb{C}^n}(\phi f)$  and the Hankel operator  $H_\phi$  from  $L_a^{2,\mathbb{C}^n}(dA_\alpha)$  into  $(L_a^{2,\mathbb{C}^n}(dA_\alpha))^\perp = L^{2,\mathbb{C}^n}(\mathbb{D}, dA_\alpha) \ominus L_a^{2,\mathbb{C}^n}(dA_\alpha)$  as  $H_\phi f = (I - P_\alpha^{\mathbb{C}^n})(\phi f)$ . For  $\phi \in L_{M_n}^\infty(\mathbb{D})$ , define  $\|\phi\|_\infty = \text{ess sup}_{z \in \mathbb{D}} \|\phi(z)\|$ . If  $\phi \in L_{M_n}^\infty(\mathbb{D})$ , then it is not difficult to see that  $\|T_\phi\| \leq \|\phi\|_\infty$  and  $\|H_\phi\| \leq \|\phi\|_\infty$ . This is so as  $\|P_\alpha^{\mathbb{C}^n}\| \leq 1$  and  $\|I - P_\alpha^{\mathbb{C}^n}\| \leq 1$ .

For  $\phi \in L_{M_n}^\infty(\mathbb{D})$ , we define the little Hankel operator  $S_\phi$  from  $L_a^{2,\mathbb{C}^n}(dA_\alpha)$  into itself as  $S_\phi f = P_\alpha^{\mathbb{C}^n} J_\alpha^n(\phi f)$  where  $J_\alpha^n : L^{2,\mathbb{C}^n}(\mathbb{D}, dA_\alpha) \rightarrow L^{2,\mathbb{C}^n}(\mathbb{D}, dA_\alpha)$  is defined as  $J_\alpha^n f(z) = f(\bar{z})$ . The map  $J_\alpha^n$  is unitary. There are also many equivalent ways of defining little Hankel operators. Let  $\overline{L_a^{2,\mathbb{C}^n}(dA_\alpha)} = \overline{L_a^2(dA_\alpha) \otimes \mathbb{C}^n}$ . For  $\phi \in L_{M_n}^\infty(\mathbb{D})$ , define  $h_\phi$  from  $L_a^{2,\mathbb{C}^n}(dA_\alpha)$  into  $\overline{L_a^{2,\mathbb{C}^n}(dA_\alpha)}$  as  $h_\phi f = \overline{P_\alpha^{\mathbb{C}^n}(\phi f)}$  where  $\overline{P_\alpha^{\mathbb{C}^n}}$  is the orthogonal projection from  $L^{2,\mathbb{C}^n}(\mathbb{D}, dA_\alpha)$  onto  $\overline{L_a^{2,\mathbb{C}^n}(dA_\alpha)}$ . It is not difficult to verify that  $h_\phi = J_\alpha^n S_\phi$ .

Let  $\mathcal{L}(H)$  be the space of all bounded linear operators from the Hilbert space  $H$  into itself. Let  $\mathcal{LC}(H)$  be the set of all compact operators in  $\mathcal{L}(H)$ . The quotient algebra  $\mathcal{L}(H)/\mathcal{LC}(H)$  is a Banach algebra called the Calkin algebra. The natural homomorphism from  $\mathcal{L}(H)$  onto  $\mathcal{L}(H)/\mathcal{LC}(H)$  is de-

noted by  $\pi$ . An operator  $T$  in  $\mathcal{L}(H)$  is a Fredholm operator if  $\pi(T)$  is an invertible element of  $\mathcal{L}(H)/\mathcal{LC}(H)$ . The collection of Fredholm operators on  $H$  is denoted by  $\mathcal{F}(H)$ . It follows from a theorem of Atkinson [1] that, an operator  $T$  in  $\mathcal{L}(H)$  is a Fredholm operator if and only if the range of  $T$  is closed,  $\dim \ker T$  is finite and  $\dim \ker T^*$  is finite. Let  $\mathcal{F}(L_a^2(dA_\alpha))$  be the set of all Fredholm operators in  $\mathcal{L}(L_a^2(dA_\alpha))$ . The classical index  $j$  is the function from  $\mathcal{F}(L_a^2(dA_\alpha))$  to  $\mathbb{Z}$  defined as  $j(T) = \dim \ker T - \dim \ker T^*$ . In this paper we have shown that if  $\phi \in (L_h^2(dA_\alpha))^\perp \cap L^\infty(\mathbb{D})$  and  $\text{Range} T_\phi^{(\alpha)}$  is closed, then  $T_\phi^{(\alpha)}$  is a Fredholm operator of index zero and  $T_\phi^{(\alpha)}$  is not of finite rank. The organization of the paper is as follows: In section 2, we establish certain preliminary results. We have shown that  $\ker h_\phi^{(\alpha)} = \{0\}$  if and only if  $\overline{\text{Range } h_\phi^{(\alpha)}} = \overline{L_a^2(dA_\alpha)}$  and  $\ker H_{fH}^{E_p^\alpha} = \{0\}$  if and only if  $\overline{\text{Range } H_{fH}^{E_p^\alpha}} = E_p^\alpha$  where  $\phi \in L^\infty(\mathbb{D})$ ,  $H, f \in L_a^2(dA_\alpha)$ . We relate the concept of common zero set and the rank of a Toeplitz operator. In section 3, we present the main results of the paper and some applications of these results. We establish that if  $\phi \in (L_h^2(dA_\alpha))^\perp \cap L^\infty(\mathbb{D})$  and  $\text{Range} T_\phi^{(\alpha)}$  is closed, then  $T_\phi^{(\alpha)}$  is a Fredholm operator of index zero and  $T_\phi^{(\alpha)}$  is not of finite rank. Further if  $\phi \in L_{M_n}^\infty(\mathbb{D})$  is such that  $T_\phi$  is Fredholm and of index zero in  $\mathcal{L}(L_a^{2, \mathbb{C}^n}(dA_\alpha))$  then there exists  $\psi \in E_{n \times n} = E \otimes M_n$  such that  $T_{\phi+\delta\psi}$  is invertible for all sufficiently small nonzero  $\delta$ .

## 2 Preliminaries

In this section we present some elementary results that will be used in establishing the main results of the paper. We relate the kernel of little Hankel operators and intermediate Hankel operators with the kernel of their adjoints. We relate the concept of common zero set and the rank of a Toeplitz operator. For  $p \geq 0$ , let  $E_p^\alpha = \overline{\text{span}}\{|z|^{2k}\bar{z}^n, k = 0, \dots, p; n = 0, 1, 2, \dots\}$ . For  $p \geq 0$ , the spaces  $E_p^\alpha$  are closed subspaces of  $L_a^2(dA_\alpha)$ . For  $\phi \in L^\infty(\mathbb{D})$ , we define the intermediate Hankel operator  $H_\phi^{E_p^\alpha} : L_a^2(dA_\alpha) \rightarrow E_p^\alpha$  by  $H_\phi^{E_p^\alpha}(f) = P_p^{(\alpha)}(\phi f)$ ,  $f \in L_a^2(dA_\alpha)$  where  $P_p^{(\alpha)}$  is the orthogonal projection from  $L^2(\mathbb{D}, dA_\alpha)$  onto  $E_p^\alpha$ . Notice that  $\overline{L_a^2(dA_\alpha)} \subseteq E_p^\alpha \subseteq ((L_a^2(dA_\alpha))_0)^\perp$  where  $(L_a^2(dA_\alpha))_0 = \{g \in L_a^2(dA_\alpha) : g(0) = 0\}$ .

**Lemma 2.1.** *Let  $\phi \in L^\infty(\mathbb{D})$ . Then  $\ker h_\phi^{(\alpha)} = \{0\}$  if and only if  $\overline{\text{Range } h_\phi^{(\alpha)}} = \overline{L_a^2(dA_\alpha)}$ .*

*Proof.* Observe that  $(S_\phi^{(\alpha)})^* = S_{\phi^+}^{(\alpha)}$  where  $\phi^+(z) = \overline{\phi(\bar{z})}$ . It is easy to see that  $f \in \ker S_\phi^{(\alpha)}$  if and only if  $f^+ \in \ker S_{\phi^+}^{(\alpha)}$ . This implies if  $\ker S_\phi^{(\alpha)} = \{0\}$  then  $\ker (S_\phi^{(\alpha)})^* = \{0\}$ . Thus  $\overline{\text{Range } S_\phi^{(\alpha)}} = L_a^2(dA_\alpha)$ . Conversely, if  $\overline{\text{Range } S_\phi^{(\alpha)}} = L_a^2(dA_\alpha)$  then  $\ker (S_\phi^{(\alpha)})^* = \{0\}$  which implies  $\ker S_\phi^{(\alpha)} = \{0\}$ .

Clearly  $h_\phi^{(\alpha)} f = 0$  for  $f \in \ker h_\phi^{(\alpha)}$ . This implies  $J_\alpha S_\phi^{(\alpha)} f = 0$  which further implies  $S_\phi^{(\alpha)} f = 0$ . Thus  $f^+ \in \ker S_{\phi^+}^{(\alpha)}$ . Hence  $J_\alpha f^+ \in \ker S_{\phi^+}^{(\alpha)} J_\alpha = \ker (h_\phi^{(\alpha)})^*$  as  $S_{\phi^+}^{(\alpha)} J_\alpha = (J_\alpha S_\phi^{(\alpha)})^* = (h_\phi^{(\alpha)})^*$ . Let  $\bar{g} \in \overline{L_a^2(dA_\alpha)}$  and  $\bar{g} \in \ker (h_\phi^{(\alpha)})^*$ . This implies  $J_\alpha \bar{g} \in \ker (S_\phi^{(\alpha)})^*$ . Hence  $(J_\alpha \bar{g})^+ \in h_\phi^{(\alpha)}$ . Therefore,  $(J_\alpha \bar{g})^+ \in h_\phi^{(\alpha)}$ . That is,  $g \in \ker h_\phi^{(\alpha)}$ . Thus if  $f \in L_a^2(dA_\alpha)$  then  $f \in \ker h_\phi^{(\alpha)}$  if and only if  $\bar{f} \in \ker (h_\phi^{(\alpha)})^*$ . Hence  $\ker h_\phi^{(\alpha)} = \{0\}$  if and only if  $\ker (h_\phi^{(\alpha)})^* = \{0\}$  and this is true if and only if  $\overline{\text{Range } h_\phi^{(\alpha)}} = \overline{L_a^2(dA_\alpha)}$ . The result follows.  $\square$

**Lemma 2.2.** *Let  $H \in L_a^2(dA_\alpha)$  and assume that  $f \in L_a^2(dA_\alpha)$  is not a polynomial. Then  $\ker H_{\bar{f}H}^{E_p^\alpha} = \{0\}$  if and only if  $\overline{\text{Range } H_{\bar{f}H}^{E_p^\alpha}} = E_p^\alpha$ .*

*Proof.* Notice that

$$\begin{aligned} \ker H_{\bar{f}H}^{*E_p^\alpha} &= \{g \in E_p^\alpha : P_\alpha(f\bar{H}g) = 0\} \\ &\supset \ker (h_{\bar{f}H}^{(\alpha)})^* \\ &= \{g \in \overline{L_a^2(dA_\alpha)} : P_\alpha(f\bar{H}g) = 0\}. \end{aligned}$$

If  $\ker H_{\bar{f}H}^{*E_p^\alpha} = \{0\}$  then  $\ker (h_{\bar{f}H}^{(\alpha)})^* = \{0\}$ . Hence  $\ker h_{\bar{f}H}^{(\alpha)} = \{0\}$ . Since for  $h \in L_a^2(dA_\alpha)$ ,  $H_{\bar{f}H}^{E_p^\alpha} h = h_{\bar{f}H}^{(\alpha)} h + P_{E_p^\alpha \ominus \overline{L_a^2(dA_\alpha)}}(\bar{f}Hh)$ , we obtain  $\ker H_{\bar{f}H}^{E_p^\alpha} = \{0\}$ .

Now suppose  $\ker H_{\bar{f}H}^{E_p^\alpha} = \{0\}$ . This implies  $\ker h_{\bar{f}H}^{(\alpha)} = \{0\}$ . Because if  $\ker h_{\bar{f}H}^{(\alpha)} \neq \{0\}$  then [5], [6], [7] there exists an inner function  $G_\alpha \in L_a^2(dA_\alpha)$  such that  $G_\alpha \in \ker h_{\bar{f}H}^{(\alpha)}$ . That is,  $h_{\bar{f}HG_\alpha}^{(\alpha)} \equiv 0$ . This is so as  $\ker h_{\bar{f}H}^{(\alpha)} = \ker S_{\bar{f}H}^{(\alpha)}$  is an invariant subspace of  $z$ . Observe that for  $\psi \in L^\infty(\mathbb{D})$ ,  $h_\psi^{(\alpha)} T_z^{(\alpha)} = h_{\psi z}^{(\alpha)}$  and  $(T_{\bar{z}}^{(\alpha)} h_\psi^{(\alpha)})^* = (h_\psi^{(\alpha)})^* T_z^{(\alpha)} = S_\psi^{(\alpha)} + J_\alpha T_z^{(\alpha)}$ . Further for  $g \in$

$L_a^2(dA_\alpha)$ ,  $S_{\psi^+}^{(\alpha)} J_\alpha T_z^{(\alpha)} g = S_{\psi^+}^{(\alpha)} (J_\alpha(zg)) = P_\alpha(J_\alpha(\psi^+ \bar{z} J_\alpha g)) = P_\alpha(\bar{\psi} z g) = (h_{\psi \bar{z}}^{(\alpha)})^* g$ . Similarly one can show that  $(T_{\bar{z}^k}^{(\alpha)} h_\psi^{(\alpha)})^* = (h_{\psi \bar{z}^k}^{(\alpha)})^*$  for all  $k = 0, 1, \dots, p$ . Thus  $h_{\bar{f} H G_\alpha}^{(\alpha)} \equiv 0$  implies  $(T_{\bar{z}^k}^{(\alpha)} h_{\bar{f} H G_\alpha}^{(\alpha)})^* \equiv 0$ . Hence  $(h_{\bar{f} H G_\alpha \bar{z}^k}^{(\alpha)})^* \equiv 0$ . This implies  $h_{\bar{f} H G_\alpha \bar{z}^k}^{(\alpha)} \equiv 0$  for all  $k = 0, 1, \dots, p$ . Hence  $\bar{f} H G_\alpha \bar{z}^k \in (\overline{L_a^2(dA_\alpha)})^\perp$ . That is,  $\langle \bar{f} H G_\alpha \bar{z}^k, \bar{z}^k \bar{g} \rangle = 0$  for all  $g \in L_a^2(dA_\alpha)$  and  $k = 0, 1, \dots, p$ . Thus  $\bar{f} H G_\alpha \in (E_p^\alpha)^\perp$ . Hence  $\ker H_{\bar{f} H}^{E_p^\alpha} \neq \{0\}$ . Thus  $\ker H_{\bar{f} H}^{E_p^\alpha} = \{0\}$  implies  $\ker h_{\bar{f} H}^{(\alpha)} = \{0\}$ . Hence  $\ker (h_{\bar{f} H}^{(\alpha)})^* = \{0\}$ . This implies  $\ker H_{\bar{f} H}^{*E_p^\alpha} = \{0\}$ . Because if  $\ker H_{\bar{f} H}^{*E_p^\alpha} \neq \{0\}$  then there exist  $0 \neq g \in E_p^\alpha \cap L^\infty$  such that  $\bar{f} H g \in (L_a^2(dA_\alpha))^\perp$ . That is,  $\bar{f} H \bar{g} \in (\overline{L_a^2(dA_\alpha)})^\perp$ . This implies  $h_{\bar{f} H \bar{g}}^{(\alpha)} \equiv 0$  and therefore  $\ker h_{\bar{f} H}^{(\alpha)} \supseteq \bar{g} L_a^2(dA_\alpha) \cap L_a^2(dA_\alpha) \neq \{0\}$ ,  $\bar{g} \in \overline{E_p^\alpha} \cap L^\infty = H^\infty$ . Thus  $\ker H_{\bar{f} H}^{E_p^\alpha} = \{0\}$  if and only if  $\ker H_{\bar{f} H}^{*E_p^\alpha} = \{0\}$ .  $\square$

If  $N$  is a subspace of  $L_a^2(dA_\alpha)$ , let  $\mathcal{Z}(N) = \{z \in \mathbb{D} : f(z) = 0 \text{ for all } f \in N\}$  which is called the zero set of functions in  $N$ . Here if  $z_1$  is a zero of multiplicity at most  $n$  of all functions in  $N$  then  $z_1$  appears  $n$  times in the set  $\mathcal{Z}(N)$  and they are treated as distinct elements of  $\mathcal{Z}(N)$ . In the following proposition, we have shown that if  $\phi \in L^\infty(\mathbb{D})$  is such that  $T_\phi^{(\alpha)}$  is a finite rank Toeplitz operator and  $\text{Card } \mathcal{Z}(\ker (T_\phi^{(\alpha)})^*) = \text{Rank of } T_\phi^{(\alpha)}$  then  $\phi \equiv 0$ . With the following result we begin to link the ideas of subspaces and zero-sets.

**Proposition 2.1.** *If  $N$  is a subspace of  $L_a^2(dA_\alpha)$  of finite codimension in  $L_a^2(dA_\alpha)$  then*

$$\mathcal{Z}(N) = \{z \in \mathbb{D} : f(z) = 0 \text{ for all } f \in N\}$$

*is a finite set.*

*Proof.* Suppose  $\mathcal{Z}(N)$  is an infinite set. Let  $\{z_j\}_{j=1}^\infty$  be distinct points of  $\mathcal{Z}(N)$  and let  $f_1, f_2, f_3, \dots, f_n$  be functions in  $L_a^2(dA_\alpha)$  such that

$$f_i(z_1) = \dots = f_i(z_{i-1}) = 0, f_i(z_i) = 1 \text{ for all } i \geq 2.$$

For example, we could take the functions  $(f_i)$  to be the polynomials. Then

$f_1, f_2, \dots$  are linearly independent modulo  $N$  i.e. if

$$\alpha_1 f_1 + \alpha_2 f_2 + \dots + \alpha_n f_n \in N$$

where  $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathbb{C}$  then  $\alpha_1 = \alpha_2 = \dots = \alpha_n = 0$ . This contradicts the assumption that  $N$  has finite codimension in  $L_a^2(dA_\alpha)$ . Since each zero of an analytic function has finite multiplicity, the result is proved.  $\square$

Let  $k_z^{1+\frac{\alpha}{2}}$  be the normalised reproducing kernel for the Bergman space  $L_a^2(dA_\alpha)$ . When  $|z| \rightarrow 1$ ,  $k_z^{1+\frac{\alpha}{2}} \rightarrow 0$  weakly and the normalised reproducing kernels  $k_z^{1+\frac{\alpha}{2}}, z \in \mathbb{D}$  span  $L_a^2(dA_\alpha)$  [13].

**Theorem 2.3.** *If  $\phi \in L^\infty(\mathbb{D})$  is such that  $T_\phi^{(\alpha)}$  is a finite rank Toeplitz operator and  $\text{Card } \mathcal{Z} \left( \ker \left( T_\phi^{(\alpha)} \right)^* \right) = \text{Rank of } T_\phi^{(\alpha)}$  then  $\phi \equiv 0$ .*

*Proof.* Suppose  $T_\phi^{(\alpha)}$  is a finite rank. Then  $\text{Range } T_\phi^{(\alpha)}$  is finite dimensional and is a closed subspace of  $L_a^2(dA_\alpha)$ . Let  $n = \dim \text{Range } T_\phi^{(\alpha)}$ . Thus  $\ker \left( T_\phi^{(\alpha)} \right)^* = \overline{(\text{Range } T_\phi^{(\alpha)})}^\perp$  has finite codimension. Therefore by Proposition 2.1,  $\mathcal{Z} \left( \ker \left( T_\phi^{(\alpha)} \right)^* \right)$ , the common zero set of  $\ker \left( T_\phi^{(\alpha)} \right)^*$  is a finite set. Without loss of generality, we shall assume the elements of  $\mathcal{Z} \left( \ker \left( T_\phi^{(\alpha)} \right)^* \right)$  are distinct. Suppose  $\mathcal{Z} \left( \ker \left( T_\phi^{(\alpha)} \right)^* \right) = \{a_1, a_2, \dots, a_n\}$ . Here  $a_1, a_2, \dots, a_n$  are all distinct. Then  $\ker \left( T_\phi^{(\alpha)} \right)^* \subset \{f \in L_a^2(dA_\alpha) : f(a_i) = 0, i = 1, 2, \dots, n\}$ . But  $\{f \in L_a^2(dA_\alpha) : f(a_i) = 0, i = 1, 2, \dots, n\} = \{f \in L_a^2(dA_\alpha) : \langle f, k_{a_i}^{1+\frac{\alpha}{2}} \rangle = 0, i = 1, 2, \dots, n\} = \left\{ k_{a_1}^{1+\frac{\alpha}{2}}, k_{a_2}^{1+\frac{\alpha}{2}}, \dots, k_{a_n}^{1+\frac{\alpha}{2}} \right\}^\perp$ . Thus  $\text{sp} \left\{ k_{a_1}^{1+\frac{\alpha}{2}}, k_{a_2}^{1+\frac{\alpha}{2}}, \dots, k_{a_n}^{1+\frac{\alpha}{2}} \right\} \subset \left( \ker \left( T_\phi^{(\alpha)} \right)^* \right)^\perp = \text{Range } T_\phi^{(\alpha)}$ . In case of repeated zeros (if  $a$  is a zero of order  $m$ , say) the derivatives of the corresponding reproducing kernel up to order  $m - 1$ , i.e.,  $k_a^{1+\frac{\alpha}{2}}, \frac{\partial}{\partial \bar{a}} k_a^{1+\frac{\alpha}{2}}, \dots, \frac{\partial^{m-1}}{\partial \bar{a}^{m-1}} k_a^{1+\frac{\alpha}{2}}$  are included in the spanned set [7]. Now since  $\text{Range } T_\phi^{(\alpha)}$  has dimension  $n$  and  $k_{a_1}^{1+\frac{\alpha}{2}}, k_{a_2}^{1+\frac{\alpha}{2}}, \dots, k_{a_n}^{1+\frac{\alpha}{2}}$  are linearly independent therefore  $\text{Range } T_\phi^{(\alpha)} = \text{sp} \left\{ k_{a_1}^{1+\frac{\alpha}{2}}, k_{a_2}^{1+\frac{\alpha}{2}}, \dots, k_{a_n}^{1+\frac{\alpha}{2}} \right\}$ . Thus  $\ker \left( T_\phi^{(\alpha)} \right)^* = \left\{ k_{a_1}^{1+\frac{\alpha}{2}}, k_{a_2}^{1+\frac{\alpha}{2}}, \dots, k_{a_n}^{1+\frac{\alpha}{2}} \right\}^\perp = \{f \in L_a^2(dA_\alpha) : f(a_i) = 0, i = 1, 2, \dots, n\}$  is an invariant subspace of the Bergman shift operator  $T_z^{(\alpha)}$  defined on  $L_a^2(dA_\alpha)$ . Since  $T_\phi^{(\alpha)}$  is finite rank implies  $\left( T_\phi^{(\alpha)} \right)^*$  is finite rank therefore one can show that

$\ker T_\phi^{(\alpha)}$  is also an invariant subspace of  $L_a^2(dA_\alpha)$ . Let  $\mathcal{P}$  be the set of all polynomials in  $L_a^2(dA_\alpha)$  and  $M = \ker T_\phi^{(\alpha)}$ . Since  $\text{Range} T_\phi^{(\alpha)}$  has dimension  $n$ , therefore  $T_\phi^{(\alpha)}1, T_\phi^{(\alpha)}z, \dots, T_\phi^{(\alpha)}z^n$  are linearly dependent. This implies there exists a non-zero polynomial  $p$  of degree at most  $n$  such that  $T_\phi^{(\alpha)}p = 0$ . That is,  $\phi p \in (L_a^2(dA_\alpha))^\perp$ . Using the facts that codimension of  $M$  is finite, and  $T_z^{(\alpha)}M \subset M$ , it follows that  $\mathcal{P} \cap M$  is a nontrivial ideal of  $\mathcal{P}$ . Since  $\mathcal{P}$  is a principal ideal ring, there exists  $q \in \mathcal{P}$  such that  $\mathcal{P} \cap M = q\mathcal{P}$ , see [8]. Thus,  $T_{\phi q}^{(\alpha)}g = 0$  for all polynomials  $g \in \mathcal{P}$ . This implies  $T_{\phi q}^{(\alpha)}z^k = 0$  for all  $k \geq 0$ . That is,  $\phi q \in (\bar{z}^k L_a^2(dA_\alpha))^\perp$  for all  $k \geq 0$ . Hence  $\phi q \in \bigcap_{k \geq 0} (\bar{z}^k L_a^2(dA_\alpha))^\perp = (\cup_{k \geq 0} \bar{z}^k L_a^2(dA_\alpha))^\perp$ . Therefore, it follows that  $\phi q \perp \bar{z}^k z^n$  for all  $k, n \geq 0$ . Now,  $\phi q \in L^\infty \subset L^2$  implies that  $\phi q = 0$ . Thus  $\phi = 0$  except at the zeros of  $q$  which is a polynomial of degree at most  $n$ . Hence  $\phi \equiv 0$  as  $\phi \in L^\infty(\mathbb{D})$ .  $\square$

For  $z$  and  $w$  in  $\mathbb{D}$ , let  $\phi_z(w) = \frac{z-w}{1-\bar{z}w}$ . These are involutive Mobius transformations on  $\mathbb{D}$ . In fact

1.  $\phi_z \circ \phi_z(w) \equiv w$ ;
2.  $\phi_z(0) = z, \phi_z(z) = 0$ ;
3.  $\phi_z$  has a unique fixed point in  $\mathbb{D}$ .

Given  $z \in \mathbb{D}$  and  $f$  any measurable function on  $\mathbb{D}$ , we define a function  $U_z f(w) = k_z^{1+\frac{\alpha}{2}}(w)f(\phi_z(w))$ . Since  $|k_z^{1+\frac{\alpha}{2}}|^2$  is the real Jacobian determinant of the mapping  $\phi_z$  (see [14]),  $U_z$  is easily seen to be a unitary operator on  $L^2(\mathbb{D}, dA_\alpha)$  and  $L_a^2(dA_\alpha)$ . It is also easy to check that  $U_z^* = U_z$ , thus  $U_z$  is a self-adjoint unitary operator. If  $\phi \in L^\infty(\mathbb{D}, dA)$  and  $z \in \mathbb{D}$  then  $U_z T_\phi^{(\alpha)} = T_{\phi \circ \phi_z}^{(\alpha)} U_z$ . This is because  $P_\alpha U_z = U_z P_\alpha$  and for  $f \in L_a^2(dA_\alpha)$ ,  $T_{\phi \circ \phi_z}^{(\alpha)} U_z f = T_{\phi \circ \phi_z}^{(\alpha)} ((f \circ \phi_z) k_z^{1+\frac{\alpha}{2}}) = P_\alpha((\phi \circ \phi_z)(f \circ \phi_z) k_z^{1+\frac{\alpha}{2}}) = P_\alpha(U_z(\phi f)) = U_z P_\alpha(\phi f) = U_z T_\phi^{(\alpha)} f$ . Let  $\text{Aut}(\mathbb{D})$  be the Lie group of all automorphisms (biholomorphic mappings) of  $\mathbb{D}$ , and  $G_0$  the isotropy subgroup at 0; i.e.,  $G_0 = \{\psi \in \text{Aut}(\mathbb{D}) : \psi(0) = 0\}$ . Notice also that  $\phi_a(z)$ , as a function in  $a$ , is one-one and onto for any fixed  $z$  in  $\mathbb{D}$ .

**Proposition 2.2.** *If  $\phi \in L^\infty(\mathbb{D})$  then  $T_\phi^{(\alpha)}$  is finite rank if and only if  $T_{\phi \circ \phi_z}^{(\alpha)}$  is finite rank. In this case  $\text{Rank of } T_\phi^{(\alpha)} = \text{Rank of } T_{\phi \circ \phi_z}^{(\alpha)}$ .*



*Proof.* Notice that  $f \in \ker T_\phi^{(\alpha)}$  if and only if  $U_z f \in \ker T_{\phi \circ \phi_z}^{(\alpha)}$ . Further since  $U_z$  is unitary,  $\dim \ker (T_\phi^{(\alpha)})^* = \dim \ker (T_{\phi \circ \phi_z}^{(\alpha)})^*$ . Thus  $\dim \text{Range } T_\phi^{(\alpha)} = \text{codim } \ker (T_\phi^{(\alpha)})^* = \text{codim } \ker (T_{\phi \circ \phi_z}^{(\alpha)})^* = \dim \text{Range } T_{\phi \circ \phi_z}^{(\alpha)}$ .  $\square$

For  $\phi \in L^\infty(\mathbb{D})$ , let  $\tilde{\phi}(z) = \langle T_\phi^{(\alpha)} k_z^{1+\frac{\alpha}{2}}, k_z^{1+\frac{\alpha}{2}} \rangle$ , the Berezin transform of  $\phi$ . It is easy to check that  $(H_\phi^{(\alpha)})^* H_\phi^{(\alpha)} = T_{|\phi|^2}^{(\alpha)} - T_\phi^{(\alpha)} T_\phi^{(\alpha)}$  (see [13]). The following also holds.

**Proposition 2.3.** For  $\phi \in L^\infty(\mathbb{D})$ ,  $MO(\phi)^2(z) = |\widetilde{|\phi|^2}(z) - |\tilde{\phi}(z)|^2 \leq \left\| H_\phi^{(\alpha)} k_z^{1+\frac{\alpha}{2}} \right\|^2 + \left\| H_{\bar{\phi}}^{(\alpha)} k_z^{1+\frac{\alpha}{2}} \right\|^2$ .

*Proof.* Observe that

$$\begin{aligned} \left\| H_\phi^{(\alpha)} k_z^{1+\frac{\alpha}{2}} \right\| &= \left\| (I - P_\alpha)(\phi k_z^{1+\frac{\alpha}{2}}) \right\| = \left\| (I - P_\alpha)U_z(\phi \circ \phi_z) \right\| \\ &= \left\| U_z(I - P_\alpha)(\phi \circ \phi_z) \right\| = \left\| (I - P_\alpha)(\phi \circ \phi_z) \right\| = \left\| \phi \circ \phi_z - P_\alpha(\phi \circ \phi_z) \right\|. \end{aligned}$$

Similarly, we have

$$\left\| H_{\bar{\phi}}^{(\alpha)} k_z^{1+\frac{\alpha}{2}} \right\| = \left\| \bar{\phi} \circ \phi_z - P_\alpha(\bar{\phi} \circ \phi_z) \right\| = \left\| \phi \circ \phi_z - \overline{P_\alpha(\bar{\phi} \circ \phi_z)} \right\|.$$

Since  $\tilde{\phi}(z) = P_\alpha(\phi \circ \phi_z)(0)$  and  $P_\alpha \bar{g}(z) = \bar{g}(0)$  for any  $g \in L_a^2(dA_\alpha)$  and all  $z \in \mathbb{D}$ , we have

$$\begin{aligned} MO(\phi)^2(z) &= |\widetilde{|\phi|^2}(z) - |\tilde{\phi}(z)|^2 \\ &= \left\| \phi \circ \phi_z - P_\alpha(\phi \circ \phi_z)(0) \right\|^2 \\ &= \left\| \phi \circ \phi_z - P_\alpha(\phi \circ \phi_z) \right\|^2 + \left\| P_\alpha(\phi \circ \phi_z) - P_\alpha(\phi \circ \phi_z)(0) \right\|^2 \\ &= \left\| H_\phi^{(\alpha)} k_z^{1+\frac{\alpha}{2}} \right\|^2 + \left\| P_\alpha(\phi \circ \phi_z) - \overline{P_\alpha(\bar{\phi} \circ \phi_z)}(0) \right\|^2 \\ &= \left\| H_\phi^{(\alpha)} k_z^{1+\frac{\alpha}{2}} \right\|^2 + \left\| P_\alpha(\phi \circ \phi_z - \overline{P_\alpha(\bar{\phi} \circ \phi_z)}) \right\|^2 \\ &\leq \left\| H_\phi^{(\alpha)} k_z^{1+\frac{\alpha}{2}} \right\|^2 + \left\| \phi \circ \phi_z - \overline{P_\alpha(\bar{\phi} \circ \phi_z)} \right\|^2 \\ &= \left\| H_\phi^{(\alpha)} k_z^{1+\frac{\alpha}{2}} \right\|^2 + \left\| H_{\bar{\phi}}^{(\alpha)} k_z^{1+\frac{\alpha}{2}} \right\|^2. \end{aligned}$$

$\square$

**Proposition 2.4.** *If  $T \in \mathcal{L}(L_a^2(dA_\alpha))$  and  $\langle Tk_z^{1+\frac{\alpha}{2}}, k_z^{1+\frac{\alpha}{2}} \rangle = 0$  for all  $z \in \mathbb{D}$  then  $T \equiv 0$ .*

*Proof.* Define  $\rho_\alpha(T)(z) = \langle Tk_z^{1+\frac{\alpha}{2}}, k_z^{1+\frac{\alpha}{2}} \rangle$  for all  $z \in \mathbb{D}$ . If  $\rho_\alpha(T) = 0$  identically, then also  $\langle TK_z^\alpha, K_z^\alpha \rangle = K_\alpha(z, z)\rho_\alpha(T)(z) = 0$  identically where  $K_z^\alpha = K_\alpha(\cdot, z)$  is the unnormalized reproducing kernel. Thus the function  $F(x, y) = \langle TK_x^\alpha, K_y^\alpha \rangle$ ; which is holomorphic in  $x$  and  $y$ , vanishes on the ‘‘anti-diagonal’’  $x = \bar{y}$ . Passing to the variables  $u, v$  defined by  $x = u + iv$ ,  $y = u - iv$ , we get a holomorphic function  $G(u, v)$  of  $u, v$ , which vanishes when  $u, v$  are real. Thus  $F(x, y) = G(u, v) \equiv 0$ . Thus even  $\langle TK_x^\alpha, K_y^\alpha \rangle = 0$  for any  $x, y$ . Since linear combinations of  $K_x^\alpha, x \in \mathbb{D}$  are dense in  $\mathcal{L}(L_a^2(dA_\alpha))$ ; it follows that  $T \equiv 0$ .  $\square$

Notice that if for all  $z \in \mathbb{D}, P_\alpha(\phi \circ \phi_z) = P_\alpha(\phi \circ \phi_z)(0) = 0$  then  $U_z T_\phi^{(\alpha)} k_z^{1+\frac{\alpha}{2}} = U_z T_\phi^{(\alpha)} U_z 1 = T_{\phi \circ \phi_z}^{(\alpha)} 1 = 0$ . Hence  $T_\phi^{(\alpha)} k_z^{1+\frac{\alpha}{2}} = 0$  for all  $z \in \mathbb{D}$  and therefore  $\langle T_\phi^{(\alpha)} k_z^{1+\frac{\alpha}{2}}, k_z^{1+\frac{\alpha}{2}} \rangle = 0$  for all  $z \in \mathbb{D}$ . By proposition 2.4,  $T_\phi^{(\alpha)} \equiv 0$  and thus  $\phi \equiv 0$ .

### 3 Fredholm Toeplitz operators of index zero

In this section we prove the main results of the paper. We have shown that if  $\phi \in (L_h^2(dA_\alpha))^\perp \cap L^\infty(\mathbb{D})$  and  $\text{Range} T_\phi^{(\alpha)}$  is closed, then  $T_\phi^{(\alpha)}$  is a Fredholm operator of index zero and  $T_\phi^{(\alpha)}$  is not of finite rank. Further we show that if  $\phi \in L_{M_n}^\infty(\mathbb{D})$  such that  $T_\phi \in \mathcal{F}(L_a^{2, \mathbb{C}^n}(dA_\alpha))$  and  $j(T_\phi) = 0$ , then there exists  $\psi \in E_{n \times n}$  such that  $T_{\phi+\delta\psi}$  is invertible for all sufficiently small nonzero  $\delta$ .

For  $\psi \in L^\infty(\mathbb{D})$ , define  $B_\psi^{(\alpha)} : L_h^2(dA_\alpha) \rightarrow (L_h^2(dA_\alpha))^\perp$  as  $B_\psi^{(\alpha)} f = (I - Q_\alpha)(\bar{\psi}f)$  where  $Q_\alpha$  is the orthogonal projection from  $L^2(\mathbb{D}, dA_\alpha)$  onto  $L_h^2(dA_\alpha)$ . The operator  $B_\psi^{(\alpha)}$  is well defined and it is easy to see that

$$Q_\alpha(\bar{z}^n z^k) = \begin{cases} \frac{n-k+1}{n+1} \bar{z}^{n-k} & \text{if } k \leq n; \\ \frac{k-n+1}{k+1} z^{k-n} & \text{if } k \geq n. \end{cases}$$

Further we can verify that  $B_{\bar{z}^n}^{(\alpha)}(z^k) \perp B_{\bar{z}^n}^{(\alpha)}(z^j)$  if  $j \neq k$  and if  $\phi = \sum_{k=s_0}^{\infty} a_k z^k$ ,  $a_{s_0} \neq 0$  (that is,  $a_0 = \dots = a_{s_0-1} = 0$ ) then  $B_\phi^{(\alpha)}(\bar{z}^n) = \sum_{k=s_0}^{\infty} a_k \cdot$

$\cdot B_{\bar{z}^n}^{(\alpha)}(z^k)$ . It is shown in [12] that if  $\Psi$  and  $\Omega$  are two functions in  $L_a^2(dA_\alpha)$  such that  $\Psi(0) = 0 = \Omega(0)$  then the operators  $B_\Psi^{(\alpha)}$  and  $B_\Omega^{(\alpha)}$  are not of finite rank in  $L_h^2(dA_\alpha)$ . In fact the set  $\left\{B_\Psi^{(\alpha)}(\bar{z}^n)\right\}_{n=1}^p$  is linearly independent for all  $p > 0$  and the set  $\left\{B_\Omega^{(\alpha)}(z^k)\right\}_{k=1}^p$  is linearly independent for all  $p > 0$ . If  $g \in L_h^2(dA_\alpha)$  and  $g = \Psi + \bar{\Omega}$ , where  $\Psi$  and  $\Omega$  are from  $L_a^2(dA_\alpha)$  then  $B_\Omega^{(\alpha)} = B_g^{(\alpha)}|_{L_a^2(dA_\alpha)}$  is of finite rank if and only if  $\Omega \equiv 0$  and similarly  $B_\Psi^{(\alpha)} = B_g^{(\alpha)}|_{\overline{L_a^2(dA_\alpha)}}$  is of finite rank if and only if  $\Psi \equiv 0$ . Notice that for  $f \in L_a^2(dA_\alpha)$ ,  $B_f^{(\alpha)}L_h^2(dA_\alpha) = B_{\bar{f}}^{(\alpha)}L_a^2(dA_\alpha)$  and we shall also write  $B_{\bar{f}}^{(\alpha)}|_{L_a^2(dA_\alpha)} = B_{\bar{f}}^{(\alpha)}$ . Thus  $B_{\bar{f}}^{(\alpha)} : L_a^2(dA_\alpha) \rightarrow (L_h^2(dA_\alpha))^\perp$  is defined as  $B_{\bar{f}}^{(\alpha)}h = (I - Q_\alpha)(\bar{f}h)$  for all  $h \in L_a^2(dA_\alpha)$  and  $\ker B_{\bar{f}}^{(\alpha)} = \{h \in L_a^2(dA_\alpha) : \bar{f}h \in L_h^2(dA_\alpha)\}$ .

From [2], it follows that if  $f$  is not constant,  $\ker B_{\bar{f}}^{(\alpha)} = \text{sp}\{1\}$ . If  $f \equiv$  constant then  $\ker B_{\bar{f}}^{(\alpha)} = L_a^2(dA_\alpha)$ , hence  $B_{\bar{f}}^{(\alpha)} \equiv 0$ . Now  $(B_{\bar{f}}^{(\alpha)})^*$  maps  $(L_h^2(dA_\alpha))^\perp$  into  $L_a^2(dA_\alpha)$  and  $(B_{\bar{f}}^{(\alpha)})^*k = P_\alpha(fk)$  for all  $k \in (L_h^2(dA_\alpha))^\perp$ . It therefore follows that for  $f \in L_a^2(dA_\alpha)$ ,

$$\overline{\text{Range} \left( B_{\bar{f}}^{(\alpha)} \right)^*} = \begin{cases} \{0\} & \text{if } f \equiv \text{constant}; \\ (\text{sp}\{1\})^\perp & \text{if } f \neq \text{constant}. \end{cases}$$

**Theorem 3.1.** *If  $\phi \in (L_h^2(dA_\alpha))^\perp \cap L^\infty(\mathbb{D})$  and  $\text{Range} T_\phi^{(\alpha)}$  is closed, then  $T_\phi^{(\alpha)}$  is a Fredholm operator of index zero and  $T_\phi^{(\alpha)}$  is not of finite rank.*

*Proof.* Given that  $T_\phi^{(\alpha)}1 = (T_\phi^{(\alpha)})^*1 = 0$ , hence  $\phi \in (L_h^2(dA_\alpha))^\perp$ . If  $\psi \in L^\infty(\mathbb{D})$  then it is not difficult to verify that  $S_\psi^{(\alpha)} \equiv 0$  if and only if  $\bar{\psi} \in (\overline{L_a^2(dA_\alpha)})^\perp$ . Thus  $S_\phi^{(\alpha)} \equiv 0$  and  $S_\phi^{(\alpha)}f = 0$  for all  $f \in L_a^2(dA_\alpha)$ . Hence  $(\bar{\phi}f) \in (\overline{L_a^2(dA_\alpha)})^\perp$  for all  $f \in L_a^2(dA_\alpha)$  and

$$\begin{aligned} \ker \left( T_\phi^{(\alpha)} \right)^* &= \left\{ f \in L_a^2(dA_\alpha) : (\bar{\phi}f) \in (L_a^2(dA_\alpha))^\perp \right\} \\ &= \left\{ f \in L_a^2(dA_\alpha) : (\bar{\phi}f) \in (L_h^2(dA_\alpha))^\perp \right\}. \end{aligned}$$

Thus

$$\begin{aligned}
\ker \left( T_{\bar{\phi}}^{(\alpha)} \right)^* &= \left\{ f \in L_a^2(dA_\alpha) : (\bar{\phi}f) \in (L_h^2(dA_\alpha))^\perp \right\} \\
&= \left\{ f \in L_a^2(dA_\alpha) : \langle \bar{\phi}f, g \rangle = 0 \text{ for all } g \in L_h^2(dA_\alpha) \right\} \\
&= \left\{ f \in L_a^2(dA_\alpha) : \langle \bar{\phi}, \bar{f}g \rangle = 0 \text{ for all } g \in L_h^2(dA_\alpha) \right\} \\
&= \left\{ f \in L_a^2(dA_\alpha) : \langle \bar{\phi}, (I - Q_\alpha)(\bar{f}g) \rangle = 0 \text{ for all } g \in L_h^2(dA_\alpha) \right\} \\
&= \left\{ f \in L_a^2(dA_\alpha) : \langle \bar{\phi}, B_{\bar{f}}^{(\alpha)} g \rangle = 0 \text{ for all } g \in L_h^2(dA_\alpha) \right\}.
\end{aligned}$$

**Case-1:** Let  $f \in L_a^2(dA_\alpha)$  be a polynomial of degree  $k \geq 1$ , and  $H(z) = z^{k+1} \in L_a^2(dA_\alpha)$ . Then  $\bar{f}H \in (E_p^\alpha)^\perp$  for all  $p \geq 0$  and  $H_{\bar{f}H}^{E_p^\alpha} \equiv 0$ . Since  $\bar{f}z^k \notin (E_p^\alpha)^\perp$ , and  $\ker H_{\bar{f}}^{E_p^\alpha}$  is an invariant subspace of  $z$ , hence  $\ker H_{\bar{f}}^{E_p^\alpha} = z^{k+1}L_a^2(dA_\alpha)$ . Therefore  $H_{\bar{f}H}^{*E_p^\alpha} \equiv 0$  and  $\ker H_{\bar{f}H}^{*E_p^\alpha} = \bar{z}^{k+1}E_p^\alpha$  for all  $p \geq 0$ .  
Now

$$\begin{aligned}
\ker \left( B_{\bar{f}}^{(\alpha)} \right)^* &= \left\{ g \in (L_h^2(dA_\alpha))^\perp : P_\alpha(fg) = 0 \right\} \\
&= \left\{ g \in (L_h^2(dA_\alpha))^\perp : fg \in (L_a^2(dA_\alpha))^\perp \right\}
\end{aligned}$$

and  $\ker H_{\bar{f}}^{*E_p^\alpha} = \{g \in E_p^\alpha : fg \in (L_a^2(dA_\alpha))^\perp\}$ . Hence  $\ker \left( B_{\bar{f}}^{(\alpha)} \right)^* \cap E_p^\alpha = \ker H_{\bar{f}}^{*E_p^\alpha} \cap (L_h^2(dA_\alpha))^\perp$  and therefore

$$\begin{aligned}
\ker \left( B_{\bar{f}}^{(\alpha)} \right)^* &= \bigcup_{p \geq 0} \left( \ker \left( B_{\bar{f}}^{(\alpha)} \right)^* \cap E_p^\alpha \right) \\
&= \bigcup_{p \geq 0} \left( \ker H_{\bar{f}}^{*E_p^\alpha} \cap (L_h^2(dA_\alpha))^\perp \right) \\
&= \left( \bigcup_{p \geq 0} \ker H_{\bar{f}}^{*E_p^\alpha} \right) \cap (L_h^2(dA_\alpha))^\perp \\
&= \left( \bigcup_{p \geq 0} \bar{z}^{k+1}E_p^\alpha \right) \cap (L_h^2(dA_\alpha))^\perp = \{0\}.
\end{aligned}$$

Thus if  $\bar{\phi} \in (L_h^2(dA_\alpha))^\perp$  and  $\bar{\phi} \neq 0$  then  $\left( B_{\bar{f}}^{(\alpha)} \right)^* \bar{\phi} \neq 0$ .

**Case-2:** If  $f \in L_a^2(dA_\alpha)$  is a constant, then  $B_{\bar{f}}^{(\alpha)} \equiv 0$  and hence  $(B_{\bar{f}}^{(\alpha)})^* \equiv 0$  and therefore  $(B_{\bar{f}}^{(\alpha)})^* \bar{\phi} = 0$  if  $\bar{\phi} \neq 0$ .

**Case-3:** If  $f \in L_a^2(dA_\alpha)$  is not a polynomial then  $\bar{f}g \notin (E_p^\alpha)^\perp$  for any  $g \in L_a^2(dA_\alpha)$ ,  $g \neq 0$ ,  $p \geq 0$ . Hence  $\ker H_{\bar{f}}^{E_p^\alpha} = \{0\}$  and therefore by Lemma 2.2, we obtain  $\ker H_{\bar{f}}^{*E_p^\alpha} = \{0\}$  for all  $p \geq 0$ . This implies  $\ker (B_{\bar{f}}^{(\alpha)})^* = \bigcup_{p \geq 0} (\ker H_{\bar{f}}^{*E_p^\alpha} \cap (L_h^2(dA_\alpha))^\perp) = \{0\}$  and  $(B_{\bar{f}}^{(\alpha)})^* \bar{\phi} \neq 0$  if  $\bar{\phi} \in (L_h^2(dA_\alpha))^\perp$  and  $\bar{\phi} \neq 0$ .

Thus from the above three cases, it follows that

$$\begin{aligned} \ker (T_\phi^{(\alpha)})^* &= \left\{ f \in L_a^2(dA_\alpha) : \langle \bar{\phi}, B_{\bar{f}}^{(\alpha)} g \rangle = 0 \text{ for all } g \in L_h^2(dA_\alpha) \right\} \\ &= \left\{ f \in L_a^2(dA_\alpha) : (B_{\bar{f}}^{(\alpha)})^* \bar{\phi} = 0 \right\} \\ &= \begin{cases} L_a^2(dA_\alpha) & \text{if } \phi \equiv 0; \\ \text{sp}\{1\} & \text{if } \phi \neq 0. \end{cases} \end{aligned}$$

Hence either  $\phi \equiv 0$  or  $T_\phi^{(\alpha)}$  is not of finite rank. □

Notice that if  $\phi \in (L_h^2(dA_\alpha))_0^\perp$  then  $P_\alpha \phi \equiv (P_\alpha \phi)(0) \equiv b$ , a constant.

**Lemma 3.2.** *Let  $\mathcal{L}$  be the vector space consisting of all square matrices of order  $k$  that are singular. Then there is a sequence of elementary operations (row and column operations) which when applied to the matrices of  $\mathcal{L}$  results in a set of matrices all of which have entry zero in the same position.*

*Proof.* We shall use mathematical induction to establish the Lemma. Assume the Lemma is valid for matrices of order  $< k$ . Notice that if  $A \in \mathcal{L}$  and  $A \neq 0$  then  $A$  can be written in the form  $A = \begin{pmatrix} I_{r \times r} & 0_{s \times s} \\ 0_{r \times r} & 0_{s \times s} \end{pmatrix}$  by elementary row and column operations where  $r + s = k$ ,  $r > 0$ ,  $s > 0$ . Without loss of generality, we shall assume to begin with that  $A \in \mathcal{L}$ . Let  $B \in \mathcal{L}$  and assume  $B = \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}$  where  $B_{11}$  and  $B_{21}$  are of order  $r$  and  $B_{12}$  and  $B_{22}$  are of orders  $s$ . Then  $\det(B + \lambda A) = 0$  for any  $\lambda \in \mathbb{C}$ . Now  $\det(B + \lambda A)$  is a polynomial in  $\lambda$  whose highest coefficient is  $\det B_{22}$ . Hence  $\det B_{22} = 0$

for any  $B \in \mathcal{L}$ . This implies that the set of lower right hand blocks of the matrices of  $\mathcal{L}$  is a vector space of singular matrices of order  $< k$ . Using our induction hypothesis, we obtain that there is a sequence of elementary operations which when applied to these blocks results in a set of matrices all of which have entry zero in the same position. The lemma follows.  $\square$

Let  $X$  be a Banach space and  $X^*$  be its dual space. A subspace  $M$  of  $X^*$  is said to be total if for every  $0 \neq x \in X$  there is an  $f \in M$  such that  $f(x) \neq 0$ . Let  $E$  be a total subspace of  $L^\infty(\mathbb{D})$  (thought as a set of linear functionals on  $L^1(\mathbb{D}, dA)$ ). Let  $E_{n \times n} = E \otimes M_n$  and  $\mathcal{A} = \{T_\phi : \phi \in E_{n \times n}\}$ .

**Theorem 3.3.** *Let  $\phi \in L^\infty_{M_n}(\mathbb{D})$  such that  $T_\phi \in \mathcal{F}(L^2_a{}^{\mathbb{C}^n}(dA_\alpha))$  and  $j(T_\phi) = 0$ . Then there exists  $\psi \in E_{n \times n}$  such that  $T_{\phi+\delta\psi}$  is invertible for all sufficiently small nonzero  $\delta$ .*

*Proof.* First assume that there exists no  $\psi \in E_{n \times n}$  such that  $T_{\phi+\delta\psi}$  is invertible for all sufficiently small nonzero  $\delta$ . We shall show then that there exists  $0 \neq f \in \ker T_\phi$  such that  $T_\psi f \in (\ker T_\phi^*)^\perp$ . Let  $T_\phi \in \mathcal{F}(L^2_a{}^{\mathbb{C}^n}(dA_\alpha))$  and  $j(T_\phi) = 0$  where  $\phi \in L^\infty_{M_n}(\mathbb{D})$ . Let  $\psi \in L^\infty_{M_n}(\mathbb{D})$ . Then  $T_{\phi+\delta\psi}$  also belongs to  $\mathcal{F}(L^2_a{}^{\mathbb{C}^n}(dA_\alpha))$  and  $j(T_{\phi+\delta\psi}) = 0$  for sufficiently small  $\delta$ . If  $T_{\phi+\delta\psi}$  is not invertible in  $\mathcal{L}(L^2_a{}^{\mathbb{C}^n}(dA_\alpha))$  then  $\ker(T_{\phi+\delta\psi}) \neq \{0\}$ . Hence for each sequence  $(\delta_n)_{n=1}^\infty$  tending to 0, there exist vectors  $f_{\delta_n}$  satisfying  $\|f_{\delta_n}\| = 1$  and  $T_{\phi+\delta_n\psi}f_{\delta_n} = 0$  for all  $n \in \mathbb{N}$ . Let  $f_{\delta_n} = g_{\delta_n} + h_{\delta_n}$ ,  $g_{\delta_n} \in \ker T_\phi$  and  $h_{\delta_n} \in (\ker T_\phi)^\perp$ . Then  $T_{\phi+\delta_n\psi}f_{\delta_n} = 0$  implies  $T_\phi h_{\delta_n} = -\delta_n T_\psi f_{\delta_n}$ . Therefore  $T_\phi h_{\delta_n} \rightarrow 0$  and by the invertibility of  $T_\phi$  as an operator from  $(\ker T_\phi)^\perp$  to  $\text{Range} T_\phi$  we obtain  $h_{\delta_n} \rightarrow 0$ . Since  $\ker T_\phi$  is finite dimensional, there is a subsequence of  $g_{\delta_n}$  which converges to a vector  $f$ . It is not difficult to see that  $f \in \ker T_\phi$  and  $\|f\| = 1$ . Moreover, since each  $T_\psi f_{\delta_n} = -\delta_n^{-1} T_\phi f_{\delta_n} \in \text{Range} T_\phi$  and  $\text{Range} T_\phi$  is closed, we have  $T_\psi f \in \text{Range} T_\phi = (\ker T_\phi^*)^\perp$ . Now assume that  $T_\phi$  is Fredholm and  $j(T_\phi) = 0$ . Assume  $\dim \ker T_\phi = \dim \ker T_\phi^* = n < \infty$ . Let  $\{f_1, f_2, \dots, f_n\}$  be a basis for  $\ker T_\phi$  and  $\{g_1, g_2, \dots, g_n\}$  be a basis for  $\ker T_\phi^*$ . Suppose the conclusion of the theorem is false. From the first part, it follows that for each  $T_\psi \in \mathcal{A}$  there is a nonzero vector  $f = \sum \alpha_i f_i \in \ker T_\phi$  such that  $\langle T_\psi f, g_j \rangle = 0$ ,  $j = 1, \dots, n$ . Hence for all  $T_\psi \in \mathcal{A}$  the  $n \times n$  matrix  $(\langle T_\psi f_i, g_j \rangle)$  is singular. Since  $\mathcal{A}$  is vector space, the set of all such matrices is also linear and Lemma 3.2 can be applied in this setting. But any set of elementary row and column operations applied to these matrices has the effect of simply

replacing the independent sets  $\{f_i\}$  and  $\{g_j\}$  by other independent sets  $\{f'_i\}$  and  $\{g'_j\}$ . Thus from Lemma 3.2 it follows that for some such pair of independent sets and for some fixed  $i$  and  $j$ ,  $\langle T_\psi f'_i, g'_j \rangle = 0$  for all  $T_\psi \in \mathcal{A}$ . But this is not possible for the following reasons. Notice that  $\mathcal{A} = \{T_\psi : \psi \in E_{n \times n}\}$ . Let  $0 \neq f = (f_1, f_2, \dots, f_n) \in L_a^{2, \mathbb{C}^n}(dA_\alpha)$  and  $0 \neq g = (g_1, g_2, \dots, g_n) \in L_a^{2, \mathbb{C}^n}(dA_\alpha)$ . Then there is a  $\psi = (\psi_{ij}) \in E_{n \times n}$  such that  $\langle T_\psi f, g \rangle \neq 0$ . Since  $P_\alpha^{\mathbb{C}^n} g = g$ , we obtain  $\langle \psi f, g \rangle \neq 0$  or  $\sum_{i,j=1}^n \psi_{ij} f_j \bar{g}_i \neq 0$ . There are indices  $i_0, j_0$  such that  $f_{j_0} \neq 0, g_{i_0} \neq 0$  as  $f$  and  $g$  are nonzero vectors in  $L_a^{2, \mathbb{C}^n}(dA_\alpha)$ . So [3] the product  $f_{j_0} \bar{g}_{i_0}$  is not almost everywhere equal to zero. Hence we can find  $\psi_{i_0 j_0}$  belonging to  $E$  such that  $\int \psi_{i_0 j_0} f_{j_0} \bar{g}_{i_0} \neq 0$ . Set  $\psi_{ij} = 0$  if  $i \neq i_0$  or  $j \neq j_0$ . This defines a  $\psi$  such that  $\langle T_\psi f, g \rangle \neq 0$ . Thus we obtain there exists  $\psi \in E_{n \times n}$  such that  $T_{\phi + \delta \psi}$  is invertible for all sufficiently small nonzero  $\delta$ .  $\square$

**Corollary 3.1.** *Let  $\phi \in C_{M_n}(\overline{\mathbb{D}}) = C(\overline{\mathbb{D}}) \otimes M_n$ . The space of invertible Toeplitz operators in  $\mathcal{T} = \{T_\phi : \phi \in C_{M_n}(\overline{\mathbb{D}})\}$  is dense in the space of Fredholm Toeplitz operators of index zero in  $\mathcal{T}$ .*

*Proof.* The proof follows from the Theorem 3.3 as  $C(\overline{\mathbb{D}})$  is a total subspace of  $L^\infty(\mathbb{D})$ .  $\square$

**Corollary 3.2.** *Let  $\phi \in (L_h^2(dA_\alpha))^\perp \cap L^\infty(\mathbb{D})$  and suppose  $\text{Range} T_\phi^{(\alpha)}$  is closed. Then either  $T_\phi^{(\alpha)}$  is invertible or  $T_\phi^{(\alpha)} = W^{(\alpha)} + F^{(\alpha)}$  where  $W^{(\alpha)} \in \mathcal{L}(L_a^2(dA_\alpha))$  is invertible and is not a Toeplitz operator and  $F^{(\alpha)}$  is a nonzero finite rank operator in  $\mathcal{L}(L_a^2(dA_\alpha))$ .*

*Proof.* From Theorem 3.1, it follows that  $T_\phi^{(\alpha)}$  is Fredholm and of index zero. Now there are two possibilities. Either  $T_\phi^{(\alpha)}$  is invertible or  $T_\phi^{(\alpha)}$  is not invertible. Suppose  $T_\phi^{(\alpha)}$  is not invertible. It follows from [11] that  $T_\phi^{(\alpha)} = W^{(\alpha)} + F^{(\alpha)}$  where  $W^{(\alpha)}$  is invertible and  $F^{(\alpha)}$  is a finite rank operator in  $\mathcal{L}(L_a^2(dA_\alpha))$ . If  $F^{(\alpha)} \equiv 0$  then  $T_\phi^{(\alpha)} = W^{(\alpha)}$  is invertible. If  $F^{(\alpha)} \neq 0$  and  $W^{(\alpha)} = T_\psi^{(\alpha)}$  for some  $\psi \in L^\infty(\mathbb{D})$ . Then  $F^{(\alpha)} = T_\phi^{(\alpha)} - T_\psi^{(\alpha)} = T_{\phi - \psi}^{(\alpha)}$  is a finite rank Toeplitz operator in  $\mathcal{L}(L_a^2(dA_\alpha))$ . From [9] it follows that  $F^{(\alpha)} \equiv 0$ . But  $F^{(\alpha)} \neq 0$ . Thus  $T_\phi^{(\alpha)} = W^{(\alpha)} + F^{(\alpha)}$  where  $F^{(\alpha)} \in \mathcal{L}(L_a^2(dA_\alpha))$  is a nonzero finite rank operator and  $W^{(\alpha)} \in \mathcal{L}(L_a^2(dA_\alpha))$  is invertible and is not a Toeplitz operator. The corollary follows.  $\square$

**Corollary 3.3.** *If for  $\phi \in L^\infty(\mathbb{D})$ ,  $T_\phi^{(\alpha)}$  is invertible and  $\psi \in C_0(\mathbb{D})$  (continuous functions on  $\mathbb{D}$  vanishing on the boundary) is such that  $T_{\phi-\psi}^{(\alpha)}$  is one-one, then  $T_{\phi-\psi}^{(\alpha)}$  is invertible.*

*Proof.* Notice that for  $\psi \in C_0(\mathbb{D})$  the Toeplitz operator  $T_\psi^{(\alpha)} \in \mathcal{L}(L_a^2(dA_\alpha))$  is compact [13]. Given that  $T_\phi^{(\alpha)}$  is invertible. The result follows from [10].  $\square$

**Corollary 3.4.** *Let  $\phi \in (L_h^2(dA_\alpha))^\perp \cap L^\infty(\mathbb{D})$ . Suppose*

$$\gamma(T_\phi^{(\alpha)}) = \inf \left\{ \|T_\phi^{(\alpha)} f\| : f \in L_a^2(dA_\alpha), \|f\| = 1, f \in \left( \ker T_\phi^{(\alpha)} \right)^\perp \right\} > 0$$

*and  $W^{(\alpha)} \in \mathcal{L}(L_a^2(dA_\alpha))$  is such that  $\|T_\phi^{(\alpha)} - W^{(\alpha)}\| < \gamma(T_\phi^{(\alpha)})$ . If  $\text{Range } W^{(\alpha)}$  is closed then  $W^{(\alpha)}$  is a Fredholm operator of index zero.*

*Proof.* Since  $\gamma(T_\phi^{(\alpha)}) > 0$ , hence it follows from [4] that  $\text{Range } T_\phi^{(\alpha)}$  is closed. From Theorem 3.1, it follows that  $T_\phi^{(\alpha)}$  is Fredholm of index zero. Now since  $\|T_\phi^{(\alpha)} - W^{(\alpha)}\| < \gamma(T_\phi^{(\alpha)})$ , it again follows from [4], that  $\dim \ker W^{(\alpha)} = \dim \ker (W^{(\alpha)})^* = 0$ . As  $\text{Range } W^{(\alpha)}$  is closed, we obtain that  $W^{(\alpha)}$  is a Fredholm operator of index zero.  $\square$

## References

- [1] Atkinson, F.V., "The normal solvability of linear equations in normed spaces", *Mat. Sb.*, **28(70)**(1951), 3-14.
- [2] Axler, S. and Zheng, D., "The Berezin transform on the Toeplitz algebra", *Studia Mathematica*, **127(2)**(1998), 113-136.
- [3] Conway, J.B., "Functions of One Complex Variable", 2nd Edition, Springer-Verlag, New York, 1978.
- [4] Conway, J.B., "A Course in Functional Analysis", 2nd Edition, Springer-Verlag, 1990.
- [5] Duren, P. L., Khavinson, D., Shapiro, H. S. and Sundberg, C., "Contractive zero-divisors in Bergman spaces", *Pacific J. Math.* **157** (1993), 37-56.



- [6] Duren, P. L., Khavinson, D., Shapiro, H. S. and Sundberg, C., “*Invariant subspaces in Bergman spaces and the biharmonic equation*”, Michigan Math. J. **41** (1994), 247-259.
- [7] Hedenmalm, H., “*A factorization theorem for square area-integrable analytic functions*”, J. Reine Angew. Math. **422**(1991), 45-68.
- [8] Herstein, I. N., “*Topics in Algebra*”, John Wiley & Sons, New York, 1975.
- [9] Luecking, D.H., “*Finite rank Toeplitz operators on the Bergman space*”, Proc. Amer. Math. soc., **136**(5)(2008), 1717-1723.
- [10] MacCluer, C.R., and Hull, A.J., “*A short proof of the Fredholm Alternative*”, International Journal of Pure and applied Mathematics, **45**(3)(2008), 379-381.
- [11] Morphy, G.J. and West, T.T., “*Decomposition of index-zero Fredholm operators*”, Proc. R. Ir. Acad., **81**(1)(1981), 49-54.
- [12] Zakariasy, L., “*The rank of Hankel operators on harmonic Bergmann spaces*”, Proc. Amer. Math. soc., **131**(4)(2002), 1177-1180.
- [13] Zhu, K., “*Operator theory in function spaces*”, Monographs and Textbooks in Pure and Applied Mathematics, Marcell Dekker, Inc. **139**, New York and Basel, 1990.
- [14] Zhu, K., “*On certain unitary operators and composition operators*”, Proceedings of Symposia in Pure Mathematics, Part 2 **51**(1990), 371-385.