

NEW CLASSES OF GENERAL TRIEQUILIBRIUM INCLUSIONS*

Muhammad Aslam Noor[†] Khalida Inayat Noor[‡]

Communicated by S. Treanță

DOI 10.56082/annalsarscimag.2026.2.179

Abstract

Some new classes of general triequilibrium inclusions are introduced and investigated. We establish the equivalence between the general triequilibrium inclusions and the fixed point problems, which is used to discuss the existence of the unique solution. Using various techniques such as resolvent methods and dynamical systems coupled with finite difference approach, we suggest and analyze a number of new multi step methods for solving triequilibrium inclusions. Convergence analysis of these methods is investigated under suitable conditions. Sensitivity analysis is also investigated. Various special cases are discussed as applications of the main results. Several open problems are suggested for future research.

Keywords: equilibrium inclusions, convex functions, fixed points, iterative methods, convergence analysis, dynamical system, sensitivity analysis.

MSC: 26D15, 26D10, 49J40, 65N35, 49J40, 90C26, 90C30.

1 Introduction

Equilibrium problems which were introduced by Blum et al. [7] and Noor et al. [55] provide us with a unified, natural, novel, innovative and general

* Accepted for publication on February 05, 2026

[†]noormaslam@gmail.com, Department of Mathematics, University of Wah, Wah Cantt, Pakistan

[‡]khalidan@gmail.com, Department of Mathematics, University of Wah, Wah Cantt, Pakistan

technique to study a wide class of problems arising in different branches of mathematical and engineering sciences. Equilibrium problems can be viewed as a novel and important generalization of the variational inequalities and variational principles. By variational principles, we mean maximum and minimum problems arising in game theory, mechanics, geometrical optics, general relativity theory, field theory, economics, transportation, differential geometry and related areas. These are fascinating interesting fields that a wide class of unrelated problems can be studied in the general and unified framework of variational inequalities and equilibrium problems. For more details of the applications and generalizations of the equilibrium problems and variational inequalities, see [5–7, 10–19, 21–23, 26–28, 28–47, 49–55, 58–65, 68] and the references therein.

One of the most difficult and important problem is the development of efficient numerical methods. Lions and Stampacchia [21] and Noor [26] proved that the quasi variational inequalities are equivalent to the fixed point problem. This alternative formulation was used to suggest and investigate three-step iterations for solving the variational inequalities. These three-step iterations contain Noor (three step) iterations [30–32], Picard method, Mann (one step) iterations and Ishikawa (two-step) iterations as special cases. Suantai et. al. [62] have also considered some novel forward-backward algorithms for optimization and their applications to compressive sensing and image inpainting. Noor iterations have influenced the research in the fixed point theory and will continue to inspire further research in fractal geometry, chaos theory, coding, number theory, spectral geometry, dynamical systems, complex analysis, nonlinear programming, graphics and computer aided design. These three-step schemes are a natural generalization of the splitting methods for solving partial differential equations.

The projected dynamical systems associated with variational inequalities were considered by Dupuis and Nagurney [13]. The novel feature of the projected dynamical system is that its set of stationary points corresponds to the set of the corresponding set of the solutions of the variational inequality problem. This dynamical system is a first order initial value problem. Consequently, equilibrium and nonlinear problems arising in various branches in pure and applied sciences can now be studied in the setting of dynamical systems. It has been shown [13, 23, 44, 47, 50, 51, 53, 54, 66] that the dynamical systems are useful in developing some efficient numerical techniques for solving variational inequalities and related optimization problems.

The sensitivity analysis provides useful information for designing or planning various equilibrium systems. Sensitivity analysis can provide new insight and stimulate new ideas and techniques for problem solving. Dafer-

mos [12] studied the sensitivity analysis of the variational inequalities using the fixed point technique. This approach has strong geometrical flavor and has been investigated for various classes of variational inequalities and their variant forms, see [12, 29, 42, 47, 53, 54, 65] and the references therein.

We would like to point out that it is not possible to establish the equivalence between the equilibrium problems and the fixed point problems. Due to these drawback, one can not suggest the multistep iterative methods for solving the equilibrium problems. To overcome its draw back and facts, Noor and Noor [24] have introduced and studied some classes of equilibrium inclusions. They have proved that the equilibrium inclusions are equivalent to the fixed point problems. These equivalent fixed point formulation have been used to suggest and analyze some classes of hybrid iterative methods. Motivated and inspired by the research activities in this direction, we introduce some new classes of extended general triequilibrium inclusions involving the maximal monotone operator. We establish the equivalence between the extended general triequilibrium inclusions and fixed point problem exploring the resolvent operator approach. This alternative equivalent formulation is used to consider the existence of the solution as well as to analyze some multi step an iterative method for solving the triequilibrium inclusions. Several special cases are discussed as applications of the triequilibrium l inclusions in Section 2. These multi step methods can be viewed as a novel generalization of the Noor (three step) iterations [25], which have applications in fixed point, fractal geometry, information technology, machine learning and medical sciences and signal processing. In section 3, we discuss the unique existence of the solution as well as to suggest several inertial iterative method along with the convergence analysis. In Section 4, dynamical system approach is applied to study the stability of the solution as well as to suggest some iterative methods for solving the extended general triequilibrium problems exploring the finite difference idea. Our results in this section can be viewed as significant refinement of the known results. Sensitivity analysis for variational inequalities has been studied by many authors using quite different techniques. In Section 5, we obtain some new results for the sensitivity analysis of the extended general triequilibrium inclusions.

One of the main purposes of this paper is to demonstrate the close connection among various classes of algorithms for the solution of the extended general equilibrium inclusions and to point out that researchers in different field of equilibrium inclusions and optimization. These results may motivate and bring a large number of novel, innovate potential applications, extensions and interesting topics in these areas. We have given only a brief introduction of this new field of triequilibrium inclusions. The interested readers

may explore this field further and discover novel and fascinating applications of the extended general equilibrium inclusions in other areas of sciences such as fractal geometry, chaos theory, coding, number theory, spectral geometry, dynamical systems, complex analysis, nonlinear programming, graphics, computer aided design and related other optimization problems. It is expected the techniques and ideas of this paper may be starting point for further research.

2 Formulations and basic facts

Let Ω be a nonempty closed convex set in a real Hilbert space \mathcal{H} . We denote by $\langle \cdot, \cdot \rangle$ and $\| \cdot \|$ be the inner product and norm, respectively. First of all, we recall some concepts from convex analysis [1, 10, 25, 54] which are needed in the derivation of the main results.

We consider the extended general triequilibrium inclusion problem. For given nonlinear operators $T, g, h = \mathcal{H} \rightarrow \mathcal{H}$, a trifunction $F(., ., .) : \mathcal{H} \times \mathcal{H} \times \mathcal{H} \rightarrow \mathcal{H}$ and maximal monotone operator $A(\cdot)$, we consider the problem of finding $\mu \in \mathcal{H}$, such that

$$0 \in \rho F(\mu, T(\mu), \nu) + g(\mu) - h(\mu) + \rho A(g(\mu)), \quad \forall \nu \in \mathcal{H}, \quad (1)$$

which is called the extended general triequilibrium inclusion.

Special Cases.

1. For $g = h$, the problem (1) reduces to finding $\mu \in \mathcal{H}$ such that

$$0 \in \rho F(\mu, T(\mu), \nu) + \rho A(g(\mu)), \quad \forall \nu \in \mathcal{H} \quad (2)$$

is known as the triequilibrium inclusion.

2. If $A(\cdot) = \partial\varphi(\cdot) : \mathcal{H} \rightarrow R \cup \{+\infty\}$, the subdifferential of a convex, proper and lower semi-continuous function $\varphi(\cdot)$, then problem (1) is equivalent to finding $\mu \in \mathcal{H}$ such that

$$\begin{aligned} & \langle \rho F(\mu, T(\mu), \nu) + g(\mu) - h(\mu), h(\nu) - g(\mu) \rangle \\ & + \rho(\varphi(h(\nu)) - \varphi(g(\mu))) \geq 0, \quad \forall \nu \in \mathcal{H}, \end{aligned} \quad (3)$$

which is called the mixed general triequilibrium variational inequality.

3. If the function $\varphi(\cdot)$ is the indicator function of a closed convex set Ω in H , that is,

$$\varphi(\mu) = \begin{cases} 0, & \text{if } \mu \in \Omega \\ +\infty, & \text{otherwise,} \end{cases}$$

then problem (3) is equivalent to finding $\mu \in \Omega$, such that

$$\langle F(\mu, T(\mu), \nu) + g(\mu) - h(\mu), h(\nu) - g(\mu) \rangle \geq 0, \quad \forall \nu \in \Omega, \quad (4)$$

is called the general triequilibrium variational inequality.

4. For $F(\mu, T(\mu), \nu) = \langle T(\mu), g(\mu) - g(\nu) \rangle$, the problem (4) reduces to finding $\mu \in \Omega$ such that

$$\langle T(\mu), g(\mu) - g(\nu) \rangle \geq 0, \quad \forall \nu \in \Omega, \quad (5)$$

is called the general variational inequality, introduced and studied by Noor [27] in 1988. For applications, modification and numerical aspects of the general variational inequalities, see [27, 32, 53, 54].

5. If $\Omega^* = \{\mu \in \mathcal{H}, \langle \mu, \nu \rangle \geq 0, \quad \forall \nu \in \Omega\}$ is a polar cone of the convex cone Ω in \mathcal{H} and $h = g$, then the problem (4) is equivalent to finding $\mu \in \mathcal{H}$, such that

$$g(\mu) \in \Omega, \quad F(\mu, T(\mu), \nu) \in \Omega^*, \quad \langle F(\mu, T(\mu), \nu), g(\mu) \rangle = 0, \quad (6)$$

is called the triequilibrium complementarity problem, which appears to be a new one. For $F(\mu, T(\mu), \nu) = T(\mu)$, the triequilibrium problem (6) reduces to finding $\mu \in \mathcal{H}$ such that

$$g(\mu) \in \Omega, \quad T(\mu) \in \Omega^*, \quad \langle T(\mu), g(\mu) \rangle = 0,$$

is known as the general complementarity problem, introduced and studied by Noor [27] in 1988, which include the nonlinear complementarity problem as a special case. For applications, formulations and generalizations of the complementarity problems, see [9, 27, 32, 41, 53, 54].

For special choices of the single valued operators $T, h, g, \mathcal{A}(\cdot, \cdot)$ the continuous trifunction $F(\cdot, \cdot, \cdot)$ and the closed convex set Ω , one can obtain a wide class of complementarity problems and variational inequality problems as special cases of the extended general triequilibrium problem (1). Thus, it is clear that the problem (1) is very general and unifying one and has numerous applications in pure and applied sciences.

We now recall some well known results and notions.

Definition 1. *If A is a set valued maximal monotone operator on \mathcal{H} , then, for a constant $\rho > 0$, the resolvent operator is defined by*

$$\mathcal{J}_A = (I + \rho A)^{-1}(\mu), \quad \forall \mu \in \mathcal{H},$$

where I is the identity operator.

It is known that the resolvent operator \mathcal{J}_A is single-valued defined on all of \mathcal{H} by Minty's theorem [58] and satisfies the following assumption.

Assumption 1. [58] *The resolvent operator J_A is nonexpansive.*

$$\|J_A(\mu) - J_A(\nu)\| \leq \|\mu - \nu\|, \quad \forall \mu, \nu \in \mathcal{H}. \quad (7)$$

Assumption 1 is used to prove the existence of a solution of extended general triequilibrium inclusions as well as in analyzing convergence of the iterative methods.

Definition 2. *The trifunction $F(.,.,.) : \mathcal{H} \times \mathcal{H} \rightarrow \mathcal{H}$ with respect to an arbitrary operator T is said to be:*

1. *Strongly jointly monotone, if there exist a constant $\alpha > 0$, such that*

$$F(\mu, T(\mu), \nu) - F(\eta, T(\eta), \nu) \geq \alpha \|\mu - \eta\|^2, \quad \forall \eta, \mu, \nu \in \mathcal{H}.$$

2. *jointly Lipschitz continuous, if there exists a constant $\beta > 0$, such that*

$$\|F(\mu, T(\mu), \nu) - F(\eta, T(\eta), \nu)\| \leq \beta \|\mu - \eta\|, \quad \forall \eta, \mu, \nu \in \mathcal{H}.$$

3. *jointly monotone, if*

$$\langle F(\mu, T(\mu), \nu) - F(\eta, T(\eta), \nu), \nu \rangle \geq 0, \quad \forall \mu, \nu \in \mathcal{H}.$$

4. *pseudo monotone, if*

$$F(\mu, T(\mu), \nu) \geq 0 \implies -F(\nu, T(\nu), \mu) \geq 0, \quad \forall \mu, \nu \in \mathcal{H}.$$

Definition 3. *An operator $T : \mathcal{H} \rightarrow \mathcal{H}$ is said to be:*

1. *Strongly monotone, if there exists a constant $\alpha > 0$, such that*

$$\langle T(\mu) - T(\nu), \mu - \nu \rangle \geq \alpha \|\mu - \nu\|^2, \quad \forall \mu, \nu \in \mathcal{H}.$$

2. Lipschitz continuous, if there exists a constant $\beta > 0$, such that

$$\|T(\mu) - T(\nu)\| \leq \beta \|\mu - \nu\|, \quad \forall \mu, \nu \in \mathcal{H}.$$

3. Monotone, if

$$\langle T(\mu) - T(\nu), \mu - \nu \rangle \geq 0, \quad \forall \mu, \nu \in \mathcal{H}.$$

4. Pseudo monotone, if

$$\langle T(\mu), \nu - \mu \rangle \geq 0 \quad \Rightarrow \quad \langle T(\nu), \nu - \mu \rangle \geq 0, \quad \forall \mu, \nu \in \mathcal{H}.$$

Remark 1. Every strongly monotone operator is a monotone operator and monotone operator is a pseudo monotone operator, but the converse is not true.

3 Resolvent methods

In this section, we use the fixed point formulation to suggest and analyze some new implicit methods for solving the extended general triequilibrium inclusions. First of all, we establish the equivalence between the extended general triequilibrium inclusions and the fixed point problem applying the resolvent operator approach.

Lemma 1. The function $\mu \in \mathcal{H}$ is a solution of the extended general triequilibrium inclusion (1), if and only if, $\mu \in \mathcal{H}$ satisfies the relation

$$g(\mu) = \mathcal{J}_A[h(\mu) - \rho F(\mu, T(\mu), \nu)], \quad \forall \nu \in \mathcal{H}, \quad (8)$$

where \mathcal{J}_A is the resolvent operator and $\rho > 0$ is a constant.

Proof. Let $\mu \in \mathcal{H}$ be a solution of (1), then, for a constant ρ and $\forall \nu \in \mathcal{H}$,

$$\begin{aligned} \rho F(\mu, T(\mu), \nu) + g(\mu) & - h(\mu) + \rho A(g(\mu)) \ni 0, \\ & \iff \\ -h(\mu) + \rho F(\mu, T(\mu), \nu) & + g(\mu) + \rho A(g(\mu)) \ni 0 \\ & \iff \\ g(\mu) & = \mathcal{J}_A[h(\mu) - \rho F(\mu, T(\mu), \nu)]. \end{aligned}$$

the required (8). □

Lemma 1 implies that the general triequilibrium inclusion (1) is equivalent to the fixed point problem (8). This equivalent fixed point formulation (8) plays an important role in deriving the main results.

From equation (8), we have

$$\mu = \mu - g(\mu) + J_A[h(\mu) - \rho F(\mu, T(\mu), \nu)].$$

We define the function Φ associated with (8) as

$$\Phi(\mu) = \mu - g(\mu) + J_A[h(\mu) - \rho F(\mu, T(\mu), \nu)]. \quad (9)$$

To prove the unique existence of the solution of the problem (1), it is enough to show that the map Φ defined by (9) has a fixed point.

Theorem 1. *Let the operator g be strongly monotone with constant $\sigma > 0$ and Lipschitz continuous with constant $\zeta > 0$, respectively. Let the bifunction $F(., ., .)$ be jointly Lipschitz continuous with constant β and the operator h be Lipschitz continuous with constant ζ_1 . If there exists a parameter $\rho > 0$, such that*

$$\rho < \frac{1-k}{\beta}, \quad k < 1, \quad \zeta_1 < 1, \quad \zeta^2 < 2\sigma, \quad (10)$$

where

$$\theta = \rho\beta + k \quad (11)$$

$$k = \sqrt{1 - 2\sigma + \zeta^2} + \zeta_1. \quad (12)$$

Then there exists a unique solution of the problem (1).

Proof. From Lemma 1, it follows that problems (8) and (1) are equivalent. Thus, it is enough to show that the map $\Phi(u)$, defined by (9) has a fixed

point. For all $\eta \neq \mu \in \mathcal{H}$, we have

$$\begin{aligned}
& \left\| \Phi(\mu) - \Phi(\eta) \right\| = \left\| \mu - \eta - (g(\mu) - g(\eta)) \right\| \\
& \quad + J_A \left\| \left[h(\mu) - \rho F(\mu, T(\mu), \nu) \right] - J_A \left[h(\eta) - \rho F(\eta, T(\eta), \nu) \right] \right\| \\
& \leq \left\| \mu - \eta - (g(\mu) - g(\eta)) \right\| \\
& \quad + \left\| h(\eta) - h(\mu) - \rho(F(\eta, T(\eta), \nu) - F(\mu, T(\mu), \nu)) \right\| \\
& \leq \left\| \mu - \eta - (g(\mu) - g(\eta)) \right\| \\
& \quad + \left\| h(\eta) - h(\mu) \right\| + \rho \left\| (F(\eta, T(\eta), \nu) - F(\mu, T(\mu), \nu)) \right\| \\
& \leq \left\| \mu - \eta - (g(\mu) - g(\eta)) \right\| + \zeta_1 \left\| \eta - \mu \right\| + \rho\beta \left\| \eta - \mu \right\|. \tag{13}
\end{aligned}$$

Since the operator g is strongly monotone with constant $\sigma > 0$ and Lipschitz continuous with constant $\zeta > 0$, it follows that

$$\begin{aligned}
\left\| \mu - \eta - (g(\mu) - g(\eta)) \right\|^2 & \leq \left\| \mu - \eta \right\|^2 - 2\langle g(\mu) - g(\eta), \mu - \eta \rangle \\
& \quad + \zeta^2 \left\| g(\mu) - g(\eta) \right\|^2 \\
& \leq (1 - 2\sigma + \zeta^2) \left\| \mu - \eta \right\|^2. \tag{14}
\end{aligned}$$

From (13) and (14), we have

$$\begin{aligned}
\left\| \Phi(\mu) - \Phi(\nu) \right\| & \leq 2 \left\{ \sqrt{(1 - 2\sigma + \zeta^2)} + \zeta_1 + \rho\beta \right\} \left\| \mu - \nu \right\| \\
& = \theta \left\| \mu - \eta \right\|,
\end{aligned}$$

where

$$\begin{aligned}
\theta & = \rho\beta + k \\
k & = 2\sqrt{1 - 2\sigma + \zeta^2} + \zeta_1.
\end{aligned}$$

From (10), it follows that $\theta < 1$, which implies that the map $\Phi(u)$ defined by (9) has a fixed point, which is the unique solution of (1). \square

The fixed point formulation (8) is applied to propose and suggest some iterative methods for solving the problem (1).

Algorithm 1. For a given μ_0 , compute the approximate solution $\{\mu_{n+1}\}$ by the iterative schemes

$$y_n = (1 - \gamma_n)\mu_n + \gamma_n\{\mu_n - g(\mu_n) + J_A[h(\mu_n) - \rho F(\mu_n, T(\mu_n), \nu)]\} \quad (15)$$

$$z_n = (1 - \beta_n)\mu_n + \beta_n\{y_n - g(y_n) + J_A[h(y_n) - \rho F(y_n, T(y_n), \nu)]\} \quad (16)$$

$$\begin{aligned} \mu_{n+1} &= (1 - \alpha_n)\mu_n \\ &+ \alpha_n\{z_n - g(z_n) + J_A[h(z_n) - \rho F(z_n, T(z_n), \nu)]\}, \end{aligned} \quad (17)$$

which are known as modified Noor iterations.

We now study the convergence analysis of Algorithm 1, which is the main motivation of our next result.

Theorem 2. Let the operators g, h and the trifunction $F(., ., .)$ satisfy all the assumptions of Theorem 1. If the condition (10) holds, then the approximate solution $\{u_n\}$ obtained from Algorithm 1 converges to the exact solution $\mu \in \mathcal{H}$ of the extended general triequilibrium inclusion (1) strongly in \mathcal{H} .

Proof. From Theorem 1, we see that there exists a unique solution $\mu \in \mathcal{H}$ of the general triequilibrium inclusions (1). Let $\mu \in H$ be the unique solution of (1). Then, using Lemma 1, we have

$$\mu = (1 - \alpha_n)\mu + \alpha_n\{\mu - g(\mu) + J_A[h(\mu) - \rho F(\mu, T(\mu), \nu)]\} \quad (18)$$

$$= (1 - \beta_n)\mu + \beta_n\{\mu - g(\mu) + J_A[h(\mu) - \rho F(\mu, T(\mu), \nu)]\}. \quad (19)$$

$$= (1 - \gamma_n)\mu + \gamma_n\{\mu - g(\mu) + J_A[h(\mu) - \rho F(\mu, T(\mu), \nu)]\}. \quad (20)$$

From (17),(18), we have

$$\begin{aligned} \|\mu_{n+1} - \mu\| &= \|(1 - \alpha_n)(\mu_n - \mu) + \alpha_n(z_n - \mu - (g(z_n) - g(\mu))) \\ &\quad + \alpha_n\{A[h(\mu_n) - \rho F(z_n, T(z_n), \nu)] - J_A[h(\mu) - \rho F(\mu, T(\mu), \nu)]\}\| \\ &\leq (1 - \alpha_n)\|\mu_n - \mu\| + \alpha_n\|z_n - \mu - (g(z_n) - g(\mu))\| \\ &\quad + \alpha_n\|h(w_n) - h(\mu) - \rho(F(z_n, T(z_n), \nu) - F(\mu, T(\mu), \nu))\| \\ &\leq (1 - \alpha_n)\|\mu_n - \mu\| + \alpha_n(k + \rho\beta)\|z_n - \mu\| \\ &= (1 - \alpha_n)\|u_n - \mu\| + \alpha_n\theta\|z_n - \mu\|, \end{aligned} \quad (21)$$

where θ is defined by (11).

In a similar way, from (19) and (16), we have

$$\begin{aligned} \|z_n - \mu\| &\leq (1 - \beta_n)\|\mu_n - \mu\| + 2\beta_n\theta\|y_n - \mu - (g(y_n) - g(\mu))\| \\ &\quad + \beta_n\|g(y_n) - g(\mu) - \rho(y_n - \mu)\| + \beta_n\eta\|y_n - \mu\| \\ &\leq (1 - \beta_n)\|\mu_n - \mu\| + \beta_n(k + \rho)\|y_n - \mu\|, \\ &\leq (1 - \beta_n)\|\mu_n - \mu\| + \beta_n\theta\|y_n - \mu\|, \end{aligned} \quad (22)$$

where θ is defined by (10).

From (15) and (19), we obtain

$$\begin{aligned} \|y_n - \mu\| &\leq (1 - \gamma_n)\|\mu_n - \mu\| + \gamma_n\theta\|\mu_n - \mu\| \\ &\leq (1 - (1 - \theta)\gamma_n)\|\mu_n - \mu\| \leq \|\mu_n - \mu\|. \end{aligned} \quad (23)$$

From (22) and (23), we obtain

$$\begin{aligned} \|z_n - \mu\| &\leq (1 - \beta_n)\|\mu_n - \mu\| + \beta_n\theta\|\mu_n - \mu\| \\ &= (1 - (1 - \theta)\beta_n)\|\mu_n - \mu\| \leq \|\mu_n - \mu\|. \end{aligned} \quad (24)$$

From the above equations, we have

$$\begin{aligned} \|\mu_{n+1} - \mu\| &\leq (1 - \alpha_n)\|\mu_n - \mu\| + \alpha_n\theta\|\mu_n - \mu\| \\ &= [1 - (1 - \theta)\alpha_n]\|\mu_n - \mu\| \\ &\leq \prod_{i=0}^n [1 - (1 - \theta)\alpha_i]\|\mu_0 - \mu\|. \end{aligned}$$

Since $\sum_{n=0}^{\infty} \alpha_n$ diverges and $1 - \theta > 0$, we have $\lim_{i \rightarrow \infty} [1 - (1 - \theta)\alpha_i] = 0$. Consequently the sequence $\{\mu_n\}$ convergence strongly to μ . From (23), and (24), it follows that the sequences $\{y_n\}$ and $\{z_n\}$ also converge to μ strongly in \mathcal{H} . This completes the proof. \square

We suggest new perturbed iterative schemes for solving the extended general triequilibrium inclusion (1).

Algorithm 2. For a given μ_0 , compute the approximate solution $\{\mu_n\}$ by the iterative schemes

$$\begin{aligned} y_n &= (1 - \gamma_n)\mu_n \\ &\quad + \gamma_n\{\mu_n - g(\mu_n) + J_A[h(\mu_n) - \rho F(\mu_n, T(\mu_n), \nu)]\} + \gamma_n h_n \\ z_n &= (1 - \beta_n)\mu_n \\ &\quad + \beta_n\{y_n - g(y_n) + J_A[h(y_n) - \rho F(y_n, T(y_n), \nu)]\} + \beta_n f_n \\ \mu_{n+1} &= (1 - \alpha_n)\mu_n \\ &\quad + \alpha_n\{z_n - g(z_n) + J_A[h(z_n) - \rho F(z_n, T(z_n), \nu)]\} + \alpha_n e_n, \end{aligned}$$

where $\{e_n\}$, $\{f_n\}$, and $\{h_n\}$ are the sequences of the elements of \mathcal{H} introduced to take into account possible inexact computations and $J_{A(\mu_n)}$ is the corresponding perturbed resolvent operator and the sequences $\{\alpha_n\}$, $\{\beta_n\}$ and $\{\gamma_n\}$ satisfy

$$0 \leq \alpha_n, \beta_n, \gamma_n \leq 1; \quad \forall n \geq 0, \quad \sum_{n=0}^{\infty} \alpha_n = \infty.$$

For $\gamma_n = 0$, we obtain the perturbed Ishikawa iterative method and for $\gamma_n = 0$ and $\beta_n = 0$, we obtain the perturbed Mann iterative schemes for solving general equilibrium inclusion (1).

Also, we can suggest the following iterative methods for solving the extended general triequilibrium inclusions.

Algorithm 3. For a given μ_0 , compute μ_{n+1} by the iterative scheme

$$\mu_{n+1} = \mu_n - g(\mu_n) + J_A[h(\mu_n) - \rho F(\mu_n, T(\mu_n), \nu)],$$

which is known as the resolvent method.

Algorithm 4. For a given μ_0 , compute μ_{n+1} by the iterative scheme

$$\mu_{n+1} = \mu_n - g(\mu_n) + J_A[h(\mu_n) - \rho F(\mu_{n+1}, T(\mu_{n+1}), \nu)],$$

which is an implicit resolvent method and is equivalent to the following two-step method.

Algorithm 5. For a given μ_0 , compute μ_{n+1} by the iterative scheme

$$\begin{aligned} z_n &= \mu_n - g(\mu_n) + J_A[h(\mu_n) - \rho F(\mu_n, T(\mu_n), \nu)] \\ \mu_{n+1} &= \mu_n - g(\mu_n) + J_A[h(\mu_n) - \rho F(z_n, T(z_n), \nu)]. \end{aligned}$$

Algorithm 6. For a given μ_0 , compute μ_{n+1} by the iterative scheme

$$\mu_{n+1} = \mu_n - g(\mu_n) + J_A[h(\mu_{n+1}) - \rho F(\mu_{n+1}, T(\mu_{n+1}), \nu)],$$

which is known as the modified resolvent method and is equivalent to the iterative method.

Algorithm 7. For a given μ_0 , compute μ_{n+1} by the iterative scheme

$$\begin{aligned} z_n &= \mu_n - g(\mu_n) + J_{A(\mu_n)}[g(\mu_n) - \rho F(\mu_n, T(\mu_n), \nu)] \\ \mu_{n+1} &= \mu_n - g(\mu_n) + J_A[h(z_n) - \rho F(z_n, T(z_n), \nu)], \end{aligned}$$

which is two-step predictor-corrector method for solving the problem (1).

We can rewrite the equation (8) as:

$$\mu = \mu - g(\mu) + J_A \left[h\left(\frac{\mu + \mu}{2}\right) - \rho F\left(\left(\frac{\mu + \mu}{2}\right), T\left(\frac{\mu + \mu}{2}\right), \nu\right) \right].$$

This fixed point formulation is used to suggest the following implicit method.

Algorithm 8. For a given μ_0 , compute μ_{n+1} by the iterative scheme

$$\begin{aligned} \mu_{n+1} &= \mu_n - g(\mu_n) \\ &+ J_A \left[h\left(\frac{\mu_n + \mu_{n+1}}{2}\right) - \rho F\left(\left(\frac{\mu_n + \mu_{n+1}}{2}\right), T\left(\frac{\mu_n + \mu_{n+1}}{2}\right), \nu\right) \right]. \end{aligned}$$

To implement the implicit method, one uses the predictor-corrector technique to obtain a new two-step method for solving the problem (1).

Algorithm 9. For a given μ_0 , compute μ_{n+1} by the iterative scheme

$$\begin{aligned} z_n &= \mu_n - g(\mu_n) + J_A[h(\mu_n) - \rho F(\mu_n, T(\mu_n), \nu)] \\ \mu_{n+1} &= \mu_n - g(\mu_n) + J_A \left[h\left(\frac{z_n + \mu_n}{2}\right) - \rho F\left(\left(\frac{z_n + \mu_n}{2}\right), T\left(\frac{z_n + \mu_n}{2}\right), \nu\right) \right]. \end{aligned}$$

We now suggest multi-step inertial methods for solving the extended general triequilibrium inclusions (1).

Algorithm 10. For given μ_0, μ_1 , compute μ_{n+1} by the recurrence relation

$$\begin{aligned} z_n &= \mu_n - \theta_n (\mu_n - \mu_{n-1}), \quad n = 1, 2, \dots \\ y_n &= (1 - \gamma_n) z_n \\ &+ \gamma_n \left\{ z_n - g(z_n) + J_A \left[h\left(\frac{z_n + \mu_n}{2}\right) - \rho F\left(\left(\frac{z_n + \mu_n}{2}\right), T\left(\frac{z_n + \mu_n}{2}\right), \nu\right) \right] \right\}, \\ t_n &= (1 - \beta_n) y_n + \beta_n \left\{ y_n - g(y_n) \right. \\ &+ J_A \left[h\left(\frac{y_n + z_n + \mu_n}{3}\right) - \rho F\left(\left(\frac{y_n + z_n + \mu_n}{3}\right), T\left(\frac{y_n + z_n + \mu_n}{3}\right), \nu\right) \right] \left. \right\}, \\ \mu_{n+1} &= (1 - \alpha_n) z_n + \alpha_n \left\{ t_n - g(t_n) + J_A \left[h\left(\frac{z_n + y_n + t_n + \mu_n}{4}\right) \right. \right. \\ &\quad \left. \left. - \rho F\left(\left(\frac{z_n + t_n + \mu_n}{4}\right), T\left(\frac{y_n + z_n + t_n + \mu_n}{4}\right), \nu\right) \right] \right\}, \end{aligned}$$

where $\alpha_n, \beta_n, \gamma_n, \theta_n \in [0, 1]$, $\forall n \geq 1$.

For $g = h$, Algorithm 10 reduces to:

Algorithm 11. For given μ_0, μ_1 , compute μ_{n+1} by the recurrence relation

$$\begin{aligned}
 z_n &= \mu_n - \theta_n (\mu_n - \mu_{n-1}), \quad n = 1, 2, \dots \dots \\
 y_n &= (1 - \gamma_n) z_n \\
 + \gamma_n &\left\{ z_n - g(z_n) + J_A \left[g\left(\frac{z_n + \mu_n}{2}\right) - \rho F\left(\left(\frac{z_n + \mu_n}{2}\right), T\left(\frac{z_n + \mu_n}{2}\right), \nu\right) \right] \right\}, \\
 t_n &= (1 - \beta_n) y_n + \beta_n \left\{ y_n - g(y_n) \right. \\
 &\quad \left. + J_A \left[g\left(\frac{y_n + z_n + \mu_n}{3}\right) - \rho F\left(\left(\frac{y_n + z_n + \mu_n}{3}\right), T\left(\frac{y_n + z_n + \mu_n}{3}\right), \nu\right) \right] \right\}, \\
 \mu_{n+1} &= (1 - \alpha_n) z_n + \alpha_n \left\{ t_n - g(t_n) \right. \\
 &\quad \left. + J_A \left[g\left(\frac{z_n + y_n + t_n + \mu_n}{4}\right) - \rho F\left(\left(\frac{z_n + t_n + \mu_n}{4}\right), T\left(\frac{y_n + z_n + t_n + \mu_n}{4}\right), \nu\right) \right] \right\},
 \end{aligned}$$

where $\alpha_n, \beta_n, \gamma_n, \theta_n \in [0, 1]$, $\forall n \geq 1$, for solving the general triequilibrium inclusions (2).

Remark 2. For different and suitable choice of the parameters ρ, η, α , operators T, g, h , the trifunction $F(., ., .)$ and convex sets, one can recover new and known iterative methods for solving general triequilibrium inclusions, equilibrium complementarity problems and related optimization problems. Using the technique and ideas of Theorem 1 and Theorem 2, one can analyze the convergence of Algorithm 10 and its special cases.

4 Dynamical systems technique

In this section, we consider the dynamical system technique for solving the extended general triequilibrium inclusions. The projected dynamical systems associated with variational inequalities were considered by Dupuis and Nagurney [13]. It is worth mentioning that the dynamical systems are the initial value and boundary value problems. Consequently, variational inequalities and nonlinear problems arising in various branches in pure and applied sciences can now be studied via the differential equations. It has been shown that the dynamical systems are useful in developing some efficient numerical techniques for solving variational inequalities and related optimization problems, see [13, 23, 32, 44, 47, 50, 52–54, 66]. We consider some

new iterative methods for solving the extended general triequilibrium inclusions and investigate the convergence analysis of these new methods involving only the monotonicity of the operators.

We now define the residue vector $R(\mu)$ by the relation

$$R(\mu) = \left\{ J_A[h(\mu) - \rho F(\mu, T(\mu), \nu)] - g(\mu) \right\}, \quad \forall \nu \in \mathcal{H}. \quad (25)$$

Invoking Lemma 1, one can easily conclude that $\mu \in \mathcal{H}$ is a solution of the problem (1), if and only if, $\mu \in \mathcal{H}$ is a zero of the equation

$$R(\mu) = 0. \quad (26)$$

We now consider a dynamical system associated with the extended general triequilibrium inclusions. Using the equivalent formulation (8), we suggest a class of resolvent dynamical systems as

$$\frac{d\mu}{dt} = \lambda \left\{ J_A[h(\mu) - \rho F(\mu, T(\mu), \nu)] - g(\mu) \right\}, \quad \mu(t_0) = \alpha, \quad (27)$$

where λ is a parameter. The system of type (27) is called the resolvent dynamical system associated with the problem (1). Here the right hand is related to the resolvent and is discontinuous on the boundary. From the definition, it is clear that the solution of the dynamical system always stays in \mathcal{H} . This implies that the qualitative results such as the existence, uniqueness and continuous dependence of the solution of (1) can be studied. The equilibrium point of the dynamical system (27) is defined as follows.

Definition 4. An element $\mu \in \mathcal{H}$, is an equilibrium point of the dynamical system (27), if,

$$\frac{d\mu}{dx} = 0.$$

Thus, it is clear that $\mu \in \mathcal{H}$ is a solution of the extended general equilibrium inclusion (1), if and only if, $\mu \in \mathcal{H}$ is an equilibrium point. This implies that $\mu \in \mathcal{H}$ is a solution of the extended general triequilibrium inclusion (1), if and only if, $\mu \in \mathcal{H}$ is an equilibrium point.

Definition 5. ([13]) The dynamical system is said to converge to the solution set S^* of (27), if, irrespective of the initial point, the trajectory of the dynamical system satisfies

$$\lim_{t \rightarrow \infty} \text{dist}(\mu(t), S^*) = 0, \quad (28)$$

where

$$\text{dist}(\mu, S^*) = \inf_{\nu \in S^*} \|\mu - \nu\|.$$

It is easy to see, if the set S^* has a unique point μ^* , then (28) implies that

$$\lim_{t \rightarrow \infty} \mu(t) = \mu^*.$$

If the dynamical system is still stable at μ^* in the Lyapunov sense, then the dynamical system is globally asymptotically stable at μ^* .

Definition 6. *The dynamical system is said to be globally exponentially stable with degree η at μ^* , if, irrespective of the initial point, the trajectory of the system satisfies*

$$\|\mu(t) - \mu^*\| \leq u_1 \|\mu(t_0) - \mu^*\| \exp(-\eta(t - t_0)), \quad \forall t \geq t_0,$$

where u_1 and η are positive constants independent of the initial point.

It is clear that the globally exponentially stability is necessarily globally asymptotically stable and the dynamical system converges arbitrarily fast.

Lemma 2. *(Gronwall Lemma) ([18]) Let $\hat{\mu}$ and $\hat{\nu}$ be real-valued nonnegative continuous functions with domain $\{t : t \leq t_0\}$ and let $\alpha(t) = \alpha_0(|t - t_0|)$, where α_0 is a monotone increasing function. If, for $t \geq t_0$,*

$$\hat{\mu} \leq \alpha(t) + \int_{t_0}^t \hat{\mu}(s) \hat{\nu}(s) ds,$$

then

$$\hat{\mu}(s) \leq \alpha(t) \exp\left\{ \int_{t_0}^t \hat{\nu}(s) ds \right\}.$$

We now establish that the trajectory of the solution of the resolvent dynamical system (27) converges to the unique solution of the extended general trifunction equilibrium inclusions (1).

Theorem 3. *Let the trifunction $F(., ., .)$ be jointly Lipschitz continuous with constant β and the operators $g, h : H \rightarrow H$ be Lipschitz continuous with constants $\zeta > 0, \zeta_1 > 0$ respectively. If $\lambda(\zeta + \zeta_1 + \rho\beta) < 1$, then, for each $\mu_0 \in \mathcal{H}$, there exists a unique continuous solution $\mu(t)$ of the dynamical system (27) with $\mu(t_0) = \mu_0$ over $[t_0, \infty)$.*

Proof. Let

$$G(\mu) = \{J_A[h(\mu) - \rho F(\mu, T(\mu), \nu)] - g(\mu)\}, \quad \forall \mu \in H,$$

where $\lambda > 0$ is a constant and $G(\mu) = \frac{d\mu}{dt}$. For $\forall \mu, \nu \in H$, we have

$$\begin{aligned} \|G(\mu) - G(\eta)\| &\leq \lambda\{J_A[h(\mu) - \rho F(\mu, T(\mu), \nu)] - J_A[h(\eta) - \rho F(\eta, T(\eta), \nu)]\| \} \\ &\quad + \lambda\|g(\mu) - g(\eta)\| \\ &= \lambda\{\|g(\mu) - g(\eta)\| + \|J_A[h(\mu) - \rho F(\mu, T(\mu), \nu)] - J_A[h(\eta) - \rho F(\eta, T(\eta), \nu)]\| \} \\ &\leq \lambda\{\|g(\mu) - g(\eta)\| + \|h(\mu) - h(\eta) - \rho(F(\mu, T(\mu), \nu) - F(\eta, T(\eta), \nu))\| \} \\ &\leq \lambda\{\|g(\mu) - g(\eta)\| + \|h(\mu) - h(\eta)\| + \rho\|F(\mu, T(\mu), \nu) - F(\eta, T(\eta), \nu)\| \} \\ &\leq \lambda\{(\zeta + \zeta_1 + \beta\rho)\|\mu - \eta\|. \end{aligned}$$

This implies that the operator $G(\mu)$ is a Lipschitz continuous with constant

$\lambda\{(\zeta + \zeta_1 + \rho\beta)\} < 1$ and for each $\mu \in \mathcal{H}$, there exists a unique and continuous solution $\mu(t)$ of the dynamical system (27), defined on an interval $t_0 \leq t < T_1$ with the initial condition $\mu(t_0) = \mu_0$. Let $[t_0, T_1)$ be its maximal interval of existence. Then we have to show that $T_1 = \infty$. Consider, for any $\mu \in \Omega(\mu)$,

$$\begin{aligned} \|G(\mu)\| &= \left\| \frac{d\mu}{dt} \right\| = \lambda\| [h(\mu) - \rho F(\mu, T(\mu), \nu)] - g(\mu) \| \\ &\leq \lambda\{ \|J_A[h(\mu) - \rho F(\mu, T(\mu), \nu)] - J_A[0]\| + \|J_A[0] - g(\mu)\| \} \\ &\leq \lambda\{\delta\{\|g(\mu) - \rho F(\mu, T(\mu), \nu)\| + \|J_A[h(\mu)] - J_A[0]\| + \|J_A[0] - g(\mu)\|\} \} \\ &\leq \lambda\{(\rho\beta + \zeta_1 + \zeta)\|u\| + 2\|J_{A(\mu)}[0]\|. \end{aligned}$$

Then

$$\begin{aligned} \|\mu(t)\| &\leq \|\mu_0\| + \int_{t_0}^t \|\mu(s)\| ds \\ &\leq (\|\mu_0\| + k_1(t - t_0)) + k_2 \int_{t_0}^t \|\mu(s)\| ds, \end{aligned}$$

where $k_1 = 2\lambda\|J_{A(\mu)}[0]\|$ and $k_2 = \delta\lambda(\rho\beta + \zeta_1 + \zeta)$. Hence, by the Gronwall Lemma 2, we have

$$\|\mu(t)\| \leq \{\|u_0\| + k_1(t - t_0)\}e^{k_2(t-t_0)}, \quad t \in [t_0, T_1).$$

This shows that the solution is bounded on $[t_0, T_1)$. So, $T_1 = \infty$. \square

Theorem 4. *If the assumptions of Theorem 3 hold, then the dynamical system (27) converges globally exponentially to the unique solution of the extended general equilibrium inclusion (1).*

Proof. Since the trifunction $F(., ., .)$ is jointly Lipschitz continuous and the operators h, g are Lipschitz continuous, it follows from Theorem 3 that the dynamical system (27) has unique solution $\mu(t)$ over $[t_0, T_1]$ for any fixed $\mu_0 \in H$. Let $\mu(t)$ be a solution of the initial value problem (27). For a given $\mu^* \in H$ satisfying (1), consider the Lyapunov function

$$L(\mu) = \lambda \|\mu(t) - \mu^*\|^2, \quad u(t) \in \mathcal{H}. \quad (29)$$

From (27) and (29), we have

$$\begin{aligned} \frac{dL}{dt} &= 2\lambda \langle \mu(t) - \mu^*, \frac{d\mu}{dt} \rangle \\ &= 2\lambda \langle \mu(t) - \mu^*, J_A[h(\mu(t)) - \rho F(\mu(t), T(\mu(t)), \nu)] - g(\mu(t)) \rangle \\ &= 2\lambda \langle \mu(t) - \mu^*, J_A[h(\mu(t)) - \rho F(\mu(t), T(\mu(t)), \nu)] - g(\mu^*) \\ &\quad + g(\mu^*) - g(\mu(t)) \rangle \\ &= -2\lambda \langle \mu(t) - \mu^*, g(\mu(t)) - g(\mu^*) \rangle \\ &\quad + 2\lambda \langle \mu(t) - \mu^*, J_A[h(\mu(t)) - \rho F(\mu(t), T(\mu(t)), \nu)] - g(\mu^*) \rangle \\ &\leq -2\lambda \langle \rho(F(\mu(t), \nu) - F(\mu^*(t), T(\mu^*(t)), \nu)), g(\mu(t)) - g(\mu^*) \rangle \\ &\quad + 2\lambda \langle \mu(t) - \mu^*(t), J_A[g(\mu(t)) - \rho F(\mu(t), T(\mu(t)), \nu)] \\ &\quad - J_A[h(\mu^*(t)) - \rho F(\mu^*(t), T(\mu^*(t)), \nu)] \rangle, \\ &\leq -2\lambda\sigma \|\mu(t) - \mu^*\|^2 + \lambda \|g(\mu(t)) - g(\mu^*)\|^2 \\ &\quad + \lambda \|J_A[h(\mu(t)) - \rho F(\mu(t), T(\mu(t)), \nu)] \\ &\quad - J_A[h(\mu^*(t)) - \rho F(\mu^*(t), T(\mu^*(t)), \nu)]\|^2. \end{aligned} \quad (30)$$

Using the jointly Lipschitz continuity of the trifunction $F(., ., .)$ and Lipschitz continuity of the operator h , we have

$$\begin{aligned} &\|J_A[h(\mu(t)) - \rho F(\mu(t), T(\mu(t)), \nu)] - J_A[h(\mu^*(t)) \\ &\quad - \rho F(\mu^*(t), T(\mu^*(t)), \nu)]\| \\ &\leq \|h(\mu(t)) - h(\mu^*(t)) - \rho(F(\mu(t), T(\mu(t)), \nu) - F(\mu^*(t), T(\mu^*(t)), \nu))\| \\ &\leq (\zeta_1 + \rho\beta) \|\mu(t) - \mu^*(t)\|. \end{aligned} \quad (32)$$

From (30) and (32), we have

$$\frac{d}{dt} \|\mu(t) - \mu^*(t)\| \leq 2\xi\lambda \|\mu(t) - \mu^*(t)\|,$$

where

$$\xi = ((\zeta_1 + \rho\beta) - 2\sigma).$$

Thus, for $\lambda = -\lambda_1$, where λ_1 is a positive constant, we have

$$\|\mu(t) - \mu^*\| \leq \|\mu(t_0) - \mu^*\| e^{-\xi\lambda_1(t-t_0)},$$

which shows that the trajectory of the solution of the dynamical system (27) converges globally exponentially to the unique solution of the extended general triequilibrium inclusions (1). \square

We use the dynamical system (27) to suggest some iterative methods for solving the extended general triequilibrium inclusion (1). These methods can be viewed in the sense of Noor [25–27] involving the double resolvent operator.

For simplicity, we take $\lambda = 1$. Thus, the dynamical system (27) becomes

$$\frac{d\mu}{dt} + g(\mu) = J_{A(\mu)}[h(\mu) - \rho F(\mu, T(\mu), \nu)], \quad \mu(t_0) = \alpha. \quad (33)$$

The forward difference scheme is used to construct the implicit iterative method. Discretizing (33), we have

$$\frac{\mu_{n+1} - \mu_n}{h_1} + g(\mu_n) = J_A[h(\mu_n) - \rho F(\mu_{n+1}, T(\mu_{n+1}), \nu)], \quad (34)$$

where h_1 is the step size.

Now, we can suggest the following implicit iterative method for solving the problem (1).

Algorithm 12. For a given μ_0 , compute μ_{n+1} by the iterative scheme

$$\mu_{n+1} = \mu_n - g(\mu_n) + J_A \left[h(\mu_n) - \rho F(\mu_{n+1}, T(\mu_{n+1}), \nu) - \frac{\mu_{n+1} - \mu_n}{h_1} \right].$$

This is an implicit method and is equivalent to the following two-step method.

Algorithm 13. For a given μ_0 , compute μ_{n+1} by the iterative scheme

$$\begin{aligned} y_n &= \mu_n - g(\mu_n) + J_A[h(\mu_n) - \rho F(\mu_n, T(\mu_n), \nu)] \\ \mu_{n+1} &= \mu_n - g(\mu_n) + J_A \left[h(\mu_n) - \rho F(y_n, T(y_n), \nu) - \frac{y_n - \mu_n}{h_1} \right]. \end{aligned}$$

Discretizing (33), we now suggest other implicit iterative method for solving the extended general triequilibrium inclusion (1)

$$\frac{\mu_{n+1} - \mu_n}{h} + g(\mu_n) = J_A[h(\mu_{n+1}) - \rho F(\mu_{n+1}, T(\mu_{n+1}), \nu)], \quad (35)$$

where h is the step size.

This formulation enables us to suggest the two-step iterative method.

Algorithm 14. For a given μ_0 , compute μ_{n+1} by the iterative scheme

$$\begin{aligned} y_n &= \mu_n - g(\mu_n) + J_A \left[h(\mu_n) - \rho F(\mu_n, T(\mu_n), \nu) \right] \\ \mu_{n+1} &= \mu_n - g(\mu_n) + J_A \left[h(y_n) - \rho F(y_n, T(y_n), \nu) - \frac{y_n - \mu_n}{h} \right]. \end{aligned}$$

Discretizing (33), we propose another implicit iterative method

$$\frac{\mu_{n+1} - \mu_n}{h_1} + g(\mu_n) = J_A \left[h(\mu_n) - \rho F(\mu_{n+1}, T(\mu_{n+1}), \nu) \right],$$

where h_1 is the step size.

For $h_1 = 1$, we can suggest an implicit iterative method for solving the problem (1).

Algorithm 15. For a given μ_0 , compute μ_{n+1} by the iterative scheme

$$\mu_{n+1} = \mu_n - g(\mu_n) + J_A \left[h(\mu_n) - \rho F(\mu_{n+1}, T(\mu_{n+1}), \nu) \right].$$

From (33), we have

$$\begin{aligned} \frac{d\mu}{dt} + g(\mu) &= J_A \left[h((1 - \alpha)\mu + \alpha\mu) \right. \\ &\quad \left. - \rho F((1 - \alpha)\mu + \alpha\mu), T((1 - \alpha)\mu + \alpha\mu), \nu) \right], \end{aligned} \quad (36)$$

where $\alpha \in [0, 1]$ is a constant.

Discretization (36) and taking $h_1 = 1$, we have

$$\begin{aligned} \mu_{n+1} &= \mu_n - g(\mu_n) + J_A \left[h((1 - \alpha)\mu_n + \alpha\mu_{n-1}), \right. \\ &\quad \left. - \rho F((1 - \alpha)\mu_n + \alpha\mu_{n+1}), T((1 - \alpha)\mu_n + \alpha\mu_{n-1}), \nu) \right], \end{aligned}$$

which is an inertial type iterative method for solving the extended general triequilibrium inclusion (1). Using the predictor-corrector techniques, we have

Algorithm 16. For a given μ_0, μ_1 , compute μ_{n+1} by the iterative schemes

$$\begin{aligned} y_n &= (1 - \alpha)\mu_n + \alpha\mu_{n-1}, \quad n = 1, 2, \dots \\ \mu_{n+1} &= \mu_n - g(\mu_n) + J_A \left[h(y_n) - \rho F(y_n, T(y_n), \nu) \right], \end{aligned}$$

which is known as the inertial two-step iterative method.

We now introduce the second order dynamical system associated with the extended general triequilibrium inclusion (1). To be more precise, we consider the problem of finding $\mu \in \mathbb{H}$ such that

$$\begin{aligned} \gamma \frac{d^2\mu}{dx^2} + \frac{d\mu}{dx} &= \lambda \left\{ J_A \left[h(\mu) - \rho F(\mu, T(\mu), \nu) \right] - g(\mu) \right\}, \quad (37) \\ \mu(a) &= \alpha, \mu(b) = \beta, \end{aligned}$$

where $\gamma > 0, \lambda > 0$ and $\rho > 0$ are constants. We would like to emphasize that the problem (37) is indeed a second order boundary value problem. In a similar way, we can define the second order initial value problem associated with the dynamical system.

The equilibrium point of the dynamical system (37) is defined as follows.

Definition 7. An element $\mu \in \mathcal{H}$, is an equilibrium point of the dynamical system (37), if,

$$\gamma \frac{d^2\mu}{dx^2} + \frac{d\mu}{dx} = 0.$$

Thus, it is clear that $\mu \in \mathcal{H}$ is a solution of the extended general triequilibrium inclusion (1), if and only if, $\mu \in \mathcal{H}$ is an equilibrium point.

From (37), we have

$$g(\mu) = J_A \left[h(\mu) - \rho F(\mu, T(\mu), \nu) \right].$$

Thus, we can rewrite (37) as follows:

$$g(\mu) = J_A \left[h(\mu) - \rho F(\mu, T(\mu), \nu) + \gamma \frac{d^2\mu}{dx^2} + \frac{d\mu}{dx} \right]. \quad (38)$$

For $\lambda = 1$, the problem (37) is equivalent to finding $\mu \in \mathcal{H}$ such that

$$\begin{aligned} \gamma \frac{d^2\mu}{dx^2} + \frac{d\mu}{dx} + g(\mu) &= J_A \left[h(\mu) - \rho F(\mu, T(\mu), \nu) \right], \quad (39) \\ \mu(a) &= \alpha, \mu(b) = \beta. \end{aligned}$$

The problem (39) is called the second dynamical system, which is in fact a second order boundary value problem. This interlink among various fields of mathematical and engineering sciences is fruitful in developing implementable numerical methods for finding the approximate solutions of the extended general triequilibrium inclusions. Consequently, one can explore the ideas and techniques of the differential equations to suggest and propose hybrid proximal point methods for solving the extended general triequilibrium variational inclusions and related optimization problems.

We discretize the second-order dynamical systems (39) using central finite difference and backward difference schemes to have

$$\begin{aligned} & \gamma \frac{\mu_{n+1} - 2\mu_n + \mu_{n-1}}{h_1^2} + \frac{\mu_n - \mu_{n-1}}{h_1} + g(\mu_n) \\ & = J_A \left[h(\mu_n) - \rho F(\mu_{n+1}, T(\mu_{n+1}), \nu) \right], \end{aligned} \quad (40)$$

where h_1 is the step size.

If $\gamma = 1, h_1 = 1$, then, from equation(40) we have

Algorithm 17. For a given μ_0 , compute μ_{n+1} by the iterative scheme

$$\mu_{n+1} = \mu_n + g(\mu_n) + J_A \left[h(\mu_n) - \rho F(\mu_{n+1}, T(\mu_{n+1}), \nu) \right],$$

which is the extraresolvent method for solving the extended general triequilibrium inclusions (1).

Algorithm 17 is an implicit method. To implement the implicit method, we use the predictor-corrector technique to suggest the method.

Algorithm 18. For given μ_0, μ_1 , compute μ_{n+1} by the iterative scheme

$$\begin{aligned} y_n &= (1 - \theta_n)\mu_n + \theta_n\mu_{n-1}, \quad n = 1, 2, \dots \\ \mu_{n+1} &= \mu_n - g(\mu_n) + J_A \left[h(\mu_n) - \rho F(y_n, T(y_n), \nu) \right], \end{aligned}$$

is called the two-step inertial iterative method, where $\theta_n \in [0, 1]$ is a constant.

In a similar way, we have the following two-step method.

Algorithm 19. For given μ_0, μ_1 , compute μ_{n+1} by the iterative scheme

$$\begin{aligned} y_n &= (1 - \theta_n)\mu_n + \theta_n\mu_{n-1}, \quad n = 1, 2, \dots \\ \mu_{n+1} &= \mu_n - g(\mu_n) + J_A \left[h(y_n) - \rho F(y_n, T(y_n), \nu) \right], \end{aligned}$$

which is also called the double inertial resolvent method for solving the extended general triequilibrium inclusions (1).

We discretize the second-order dynamical systems (27) using central finite difference and backward difference schemes to have

$$\begin{aligned} & \gamma \frac{\mu_{n+1} - 2\mu_n + \mu_{n-1}}{h_1^2} + \frac{\mu_n - \mu_{n-1}}{h_1} + g(\mu_n) \\ & = J_A \left[h(\mu_n) - \rho F(\mu_{n+1}, T(\mu_{n+1}), \nu) \right], \end{aligned}$$

where h_1 is the step size.

Using this discrete form, we can suggest the following an iterative method for solving the extended general triequilibrium inclusions (1).

Algorithm 20. For given μ_0, μ_1 , compute μ_{n+1} by the iterative scheme

$$\begin{aligned} & \mu_{n+1} = \mu_n - g(\mu_n) \\ & + J_A \left[h(\mu_{n+1}) - \rho F(\mu_{n+1}, T(\mu_{n+1}), \nu) - \gamma \frac{\mu_{n+1} - 2\mu_n + \mu_{n-1}}{h_1^2} + \frac{\mu_n - \mu_{n-1}}{h_1} \right], \\ & n = 1, 2, \dots \end{aligned}$$

Algorithm 20 is called the hybrid inertial proximal method for solving the extended general triequilibrium inclusions and related optimization problems. This is a new proposed method.

Note that, for $\gamma = 1, h_1 = 1$, Algorithm 20 reduces to the following iterative method.

Algorithm 21. For given μ_0 , compute μ_{n+1} by the iterative scheme

$$\begin{aligned} \mu_{n+1} & = \mu_n - g(\mu_n) \\ & + J_A \left[h(\mu_{n+1}) + \mu_{n+1} - \mu_n - \rho F(\mu_{n+1}, T(\mu_{n+1}), \nu) \right], \end{aligned}$$

which is called the resolvent method.

We now consider the third order dynamical systems associated with the extended general triequilibrium inclusions of the type (1). To be more precise, we consider the problem of finding $\mu \in \mathcal{H}$, such that

$$\begin{aligned} \gamma \frac{d^3 \mu}{dt^3} + \zeta \frac{d^2 \mu}{dt^2} + \xi \frac{d\mu}{dt} + g(\mu) & = J_A [h(\mu) - \rho F(\mu, T(\mu), \nu)], \quad (41) \\ \text{Boundary Conditions} \quad u(a) = \alpha, \dot{\mu}(a) = \beta, \dot{\mu}(b) = \beta_1, & \end{aligned}$$

where $\gamma > 0, \zeta, \xi, \beta, \alpha, \beta_1$ and $\rho > 0$ are constants. Problem (41) is called third order dynamical system associated with extended general triequilibrium inclusions (1). The equilibrium point of the dynamical system (41) is defined as follows.

Definition 8. An element $\mu \in \mathcal{H}$, is an equilibrium point of the dynamical system (37) if

$$\gamma \frac{d^3 \mu}{dt^3} + \zeta \frac{d^2 \mu}{dt^2} + \xi \frac{d\mu}{dt} = 0.$$

Thus, it is clear that $\mu \in \mathcal{H}$ is a solution of the general equilibrium inclusion (1), if and only if, $\mu \in \mathcal{H}$ is an equilibrium point.

Consequently, the problem (27) can be written as

$$g(\mu) = J_A \left[h(\mu) - \rho F(\mu, T(\mu), \nu) + \gamma \frac{d^3 \mu}{dt^3} + \zeta \frac{d^2 \mu}{dt^2} + \xi \frac{d\mu}{dt} \right]. \quad (42)$$

We discretize the third-order dynamical systems (41) using central finite difference and backward difference schemes to have

Algorithm 22. For given μ_0, μ_1, μ_2 , compute μ_{n+1} by the iterative scheme

$$\begin{aligned} & \gamma \frac{\mu_{n+2} - 2\mu_{n+1} + 2\mu_{n-1} - \mu_{n-2}}{2h_1^3} + \zeta \frac{\mu_{n+1} - 2\mu_n + \mu_{n-1}}{h_1^2} \\ & + \xi \frac{3\mu_n - 4\mu_{n-1} + \mu_{n-2}}{2h_1} + g(\mu_n) = J_A \left[h(\mu_n) - \rho F(\mu_{n+1}, T(\mu_{n+1}), \nu) \right], \\ & n = 1, 2, \dots, \end{aligned}$$

where h_1 is the step size.

Similarly, discretizing dynamical systems (42) using central finite difference and backward difference schemes, we have

$$\begin{aligned} \mu_{n+1} = \mu_n - g(\mu_n) + J_A \left[\left\{ h(\mu_n) - \rho F(\mu_{n+1}, T(\mu_{n+1}), \nu) \right\} \right. \\ \left. + \gamma \frac{\mu_{n+2} - 2\mu_{n+1} + 2\mu_{n-1} - \mu_{n-2}}{2h_1^3} + \zeta \frac{\mu_{n+1} - 2\mu_n + \mu_{n-1}}{h_1^2} \right. \\ \left. + \xi \frac{3\mu_n - 4\mu_{n-1} + \mu_{n-2}}{2h_1} \right]. \quad (43) \end{aligned}$$

If $\gamma = 1, h_1 = 1, \zeta = 1, \xi = 1$, then, from equation(43) after adjustment, we have

Algorithm 23. For a given μ_0 , compute u_{n+1} by the iterative scheme

$$u_{n+1} = \mu_n - g(\mu_n) + J_A[h(\mu_n) - \rho F(\mu_{n+1}, T(\mu_{n+1}), \nu) + \frac{\mu_{n+1} + 3\mu_n}{2}].$$

which is an inertial type hybrid iterative methods for solving the extended general triequilibrium inclusions (1).

Remark 3. For appropriate and suitable choice of the operators T, g, h , the bifunction $F(., ., .)$, convex set, parameters and the spaces, one can suggest a wide class of implicit, explicit and inertial type methods for solving extended general trifunction equilibrium inclusions and related optimization problems.

5 Sensitivity analysis

In recent years variational inequalities are being used as mathematical programming models to study a large number of equilibrium problems arising in finance, economics, transportation, operations research and engineering sciences. The behavior of such problems as a result of changes in the problem data is always of concern, which is called sensitivity analysis. Dafermos [12] considered the sensitivity analysis considered the sensitivity of the variational inequalities using essentially the projection method. These results were extended for variational inequalities by Noor [29] and for variational inclusions by Noor et al. [42, 53, 54]. We like to mention that sensitivity analysis is important for several reasons. First, estimating problem data often introduces measurement errors, sensitivity analysis helps in identifying sensitive parameters that should be obtained with relatively high accuracy. Second, sensitivity analysis may help to predict the future changes of the equilibrium as a result of changes in the governing system. Third, sensitivity analysis provides useful information for designing or planning various equilibrium systems. Furthermore, from mathematical and engineering point of view, sensitivity analysis can provide new insight regarding problems being studied can stimulate new ideas and techniques for problem solving the problems due to these and other reasons. In this section, we study the sensitivity analysis of the extended general triequilibrium inclusions, that is, examining how solutions of such problems change when the data of the problems are changed.

We now consider the parametric versions of the problem (1). To formulate the problem, let M be an open subset of \mathcal{H} in which the parameter λ takes values. Let $g(\mu, \lambda)$ be given identity operator defined on $\mathcal{H} \times \mathcal{H} \times M$

and take value in $\mathcal{H} \times \mathcal{H}$. From now onward, we denote $g_\lambda(\cdot) \equiv g(\cdot, \lambda)$ and $F_\lambda(\cdot) \equiv F(\cdot, \lambda)$, respectively, unless otherwise specified.

The parametric extended general triequilibrium inclusions problem is to find $\mu \in \mathcal{H}$ such that

$$0 \in \rho F_\lambda(\mu, T_\lambda(\mu), \nu) + g_\lambda(\mu) - h_\lambda(\mu) + \rho A(g_\lambda(\mu)), \quad \forall \nu \in \mathcal{H} \times M. \quad (44)$$

We also assume that, for some $\bar{\lambda} \in M$, the problem (44) has a unique solution $\bar{\mu}$. From Lemma 1, we see that the parametric extended general triequilibrium inclusion are equivalent to the fixed point problem:

$$g_\lambda(\mu) = J_A \left[h_\lambda(\mu) - \rho F_\lambda(\mu, T_\lambda(\mu), \nu) \right],$$

or equivalently

$$\mu = \mu - g_\lambda(\mu) + J_A[h_\lambda(\mu) - \rho F_\lambda(\mu, T_\lambda(\mu), \nu)].$$

We now define the mapping Φ_λ associated with the problem (44) as

$$\begin{aligned} \Phi_\lambda(\mu) &= \mu - g_\lambda(\mu) \\ &+ J_A[h_\lambda(\mu) - \rho F_\lambda(\mu, T_\lambda(\mu), \nu)], \quad \forall (\mu, \lambda) \in \mathcal{H} \times M. \end{aligned} \quad (45)$$

We use this equivalence to study the sensitivity analysis of the extended general triequilibrium inclusion. We assume that for some $\bar{\lambda} \in M$, problem (44) has a solution $\bar{\mu}$ and X is a closure of a ball in \mathcal{H} centered at $\bar{\mu}$. We want to investigate those conditions under which, for each λ in a neighborhood of $\bar{\lambda}$, problem (44) has a unique solution $\mu(\lambda)$ near $\bar{\mu}$ and the function $u(\lambda)$ is (Lipschitz) continuous and differentiable.

Definition 9. Let $F_\lambda(\cdot)$ be a trifunction on $X \times M$. Then, the trifunction $F_\lambda(\cdot, \cdot, \cdot)$ with respect to an arbitrary operator T_λ is said to be:

(a) *Locally strongly jointly monotone with constant $\sigma > 0$, if*

$$\langle F_\lambda(\mu, T_\lambda(\mu), \nu) - F_\lambda(\eta, T_\lambda(\eta), \nu), \nu \rangle \geq \sigma \|\mu - \eta\|^2, \quad \forall \lambda \in M, \eta, \mu, \nu \in X.$$

(b) *jointly locally Lipschitz continuous with constant $\zeta > 0$, if*

$$\|F_\lambda(\mu, T_\lambda(\mu), \nu) - F_\lambda(\eta, T_\lambda(\eta), \nu)\| \leq \zeta \|\mu - \eta\|, \quad \forall \lambda \in M, \eta, \mu, \nu \in X.$$

Definition 10. An operator $T_\lambda : \mathcal{H} \rightarrow \mathcal{H}$ is said to be:

1. *locally strongly monotone, if there exists a constant $\alpha > 0$, such that*

$$\langle T_\lambda(\mu) - T_\lambda(\nu), \mu - \nu \rangle \geq \alpha \|\mu - \nu\|^2, \quad \forall \mu, \nu \in \mathcal{H}.$$

2. *locally Lipschitz continuous, if there exists a constant $\beta > 0$, such that*

$$\|T_\lambda(\mu) - T_\lambda(\nu)\| \leq \beta \|\mu - \nu\|, \quad \forall \mu, \nu \in \mathcal{H}.$$

3. *locally monotone, if*

$$\langle T_\lambda(\mu) - T_\lambda(\nu), \mu - \nu \rangle \geq 0, \quad \forall \mu, \nu \in \mathcal{H}.$$

We consider the case, when the solutions of the parametric extended general triequilibrium inclusion (44) lie in the interior of X . Following the ideas of Dafermos [13], Noor [29] and Noor et al. [42], we consider the map $\Phi_\lambda(\mu)$ as defined by (45). We have to show that the map $\Phi_\lambda(\mu)$ has a fixed point, which is a solution of the parametric extended general triequilibrium inclusion (44). First of all, we prove that the map $\Phi_\lambda(\mu)$, defined by (45), is a contraction map with respect to μ uniformly in $\lambda \in M$.

Lemma 3. *Let $g_\lambda(\cdot)$ be a locally strongly monotone with constants $\sigma > 0$ and locally Lipschitz continuous with constants $\zeta > 0$, respectively, and the Assumption 1 hold. If the trifunction $F_\lambda(\cdot, \cdot, \cdot)$ is locally jointly Lipschitz continuous and operator with constant β and h_λ be locally Lipschitz continuous with constant ζ_1 , we have*

$$\|\Phi_\lambda(\mu_1) - \Phi_\lambda(\mu_2)\| \leq \theta \|\mu_1 - \mu_2\|,$$

for

$$\rho < \frac{1-k}{\beta} \quad k < 1, \quad \zeta_1 < 1, \quad \zeta^2 < 2\sigma, \tag{46}$$

where

$$\theta = \left\{ \sqrt{1 - 2\sigma + \zeta^2} + \zeta_1 + \rho\beta \right\} = \{k + \rho\beta\} \tag{47}$$

and

$$k = \sqrt{1 - 2\sigma + \zeta^2} + \zeta_1. \tag{48}$$

Proof. In order to prove the existence of a solution of (44), it is enough to show that the mapping $\Phi_\lambda(\mu)$, defined by (45), is a contraction mapping.

For $\mu_1 \neq \mu_2 \in \mathcal{H}$, and using the Assumption 1, we have

$$\begin{aligned}
 & \|\Phi_\lambda(\mu_1) - \Phi_\lambda(\mu_2)\| \leq \|\mu_1 - \mu_2 - (g_\lambda(\mu_1) - g_\lambda(\mu_2))\| \\
 & + \|J_A[h_\lambda(\mu_1) - \rho F_\lambda(\mu_1, T_\lambda(\mu_1), \nu)] - J_A[h_\lambda(\mu_2) - \rho F_\lambda(\mu_2, T_\lambda(\mu_2), \nu)]\| \\
 & \leq \|\mu_1 - \mu_2 - (g_\lambda(\mu_1) - g_\lambda(\mu_2))\| \\
 & + \|h_\lambda(\mu_1) - h_\lambda(\mu_2) - \rho(F_\lambda(\mu_1, T_\lambda(\mu_1), \nu) - F_\lambda(\mu_2, T_\lambda(\mu_2), \nu))\| \\
 & \leq \|\mu_1 - \mu_2 - (g_\lambda(\mu_1) - g_\lambda(\mu_2))\| + \rho\|F_\lambda(\mu_1, T_\lambda(\mu_1), \nu) - F_\lambda(\mu_2, T_\lambda(\mu_2), \nu)\| \\
 & + \|h_\lambda(\mu_1) - h_\lambda(\mu_2)\| + \rho\|F_\lambda(\mu_1, T_\lambda(\mu_1), \nu) - F_\lambda(\mu_2, T_\lambda(\mu_2), \nu)\| \\
 & \leq \|\mu_1 - \mu_2 - (g_\lambda(\mu_1) - g_\lambda(\mu_2))\| + \zeta_1\|\mu - \nu\| + \rho\beta\|\mu_1 - \mu_2\|. \quad (49)
 \end{aligned}$$

Since the operator g_λ is a locally strongly monotone with constant $\sigma > 0$ and locally Lipschitz continuous with constant $\zeta > 0$, it follows that

$$\begin{aligned}
 \|\mu_1 - \mu_2 - (g_\lambda(\mu_1) - g_\lambda(\mu_2))\|^2 & \leq \|u_1 - u_2\|^2 - 2\langle g_\lambda(\mu_1) - g_\lambda(\mu_2), \mu_1 - \mu_2 \rangle \\
 & \quad + \|g_\lambda(\mu_1) - g_\lambda(\mu_2)\|^2 \\
 & \leq (1 - 2\sigma + \zeta^2)\|\mu_1 - \mu_2\|^2. \quad (50)
 \end{aligned}$$

From (48), (49), (50) and using the locally Lipschitz continuity of the bi-function $F(\cdot, \cdot)$ and the operator h_λ , we have

$$\begin{aligned}
 \|\Phi_\lambda(\mu_1) - \Phi_\lambda(\mu_2)\| & \leq \left\{ \zeta_1 + \sqrt{(1 - 2\sigma + \zeta^2)} + \rho\beta \right\} \|\mu_1 - \mu_2\| \\
 & = \theta\|\mu_1 - \mu_2\|,
 \end{aligned}$$

where

$$\theta = k + \rho\beta.$$

From (46), it follows that $\theta < 1$. Thus it follows that the mapping $\Phi_\lambda(\mu)$, defined by (45), is a contraction mapping and consequently it has a fixed point, which belongs to \mathcal{H} satisfying extended quasi general triequilibrium inclusion (44), the required result. \square

Remark 4. From Lemma 3, we see that the map $\Phi_\lambda(\mu)$ defined by (45) has a unique fixed point $\mu(\lambda)$, that is, $\mu(\lambda) = \Phi_\lambda(\mu)$. Also, by the assumption, the function $\bar{\mu}$, for $\lambda = \bar{\lambda}$ is a solution of the parametric extended general triequilibrium inclusion (44). Again using Lemma 3, we see that $\bar{\mu}$, for $\lambda = \bar{\lambda}$, is a fixed point of $\Phi_\lambda(\mu)$ and it is also a fixed point of $\Phi_{\bar{\lambda}}(\mu)$. Consequently, we conclude that

$$\mu(\bar{\lambda}) = \bar{\mu} = \Phi_{\bar{\lambda}}(\mu(\bar{\lambda})).$$

Using Lemma 3, we can prove the continuity of the solution $\mu(\lambda)$ of the parametric general triequilibrium inclusion (44) using the technique of Noor [46].

Lemma 4. *Assume that the triffunffion F_λ and the operator h_λ are locally Lipschitz continuous with respect to the parameter λ . If the operator $g_\lambda(\cdot)$ is locally Lipschitz continuous and the map $\lambda \rightarrow J_A$ is continuous (or Lipschitz continuous), then the function $u(\lambda)$ satisfying (45) is (Lipschitz) continuous at $\lambda = \bar{\lambda}$.*

We now state and prove the main result of this paper and is the motivation our next result.

Theorem 5. *Let $\bar{\mu}$ be the solution of the parametric extended general triequilibrium inclusion (44) for $\lambda = \bar{\lambda}$. Let the triffunffion $F_\lambda(\cdot, \cdot, \cdot)$ be jointly locally Lipschitz continuous and the operator $h_\lambda(\mu)$ be the locally strongly monotone Lipschitz continuous operator for all $\mu, \nu \in X$. If the map $\lambda \rightarrow J_{A_\mu}$ is (Lipschitz) continuous at $\lambda = \bar{\lambda}$ and the operator g_λ is locally strongly monotone Lipschitz continuous, then there exists a neighborhood $N \subset M$ of $\bar{\lambda}$ such that for $\lambda \in N$, the parametric general triffunffion equilibrium inclusion (45) has a unique solution $\mu(\lambda)$ in the interior of X , $u(\bar{\lambda}) = \bar{u}$ and $u(\lambda)$ is (Lipschitz) continuous at $\lambda = \bar{\lambda}$.*

Proof. Its proof follows from Lemma 3, Lemma 4 and Remark 4. □

6 Conclusion

Some new classes of extended general triequilibrium inclusions are introduced and investigated. We have proved that the extended general triequilibrium inclusions are equivalent to the fixed point problem. The equivalence between the general triequilibrium inclusions and fixed point problems is used to suggest some new multi step multi-step iterative methods for solving the general triffunffion equilibrium inclusions. These new methods include extra resolvent multi step hybrid resolvent methods as special cases. Convergence analysis of the proposed method is discussed for strongly monotone and Lipschitz continuous operators. Sensitivity analysis is also investigated for general triffunffion equilibrium inclusions using the equivalent fixed point approach. Iterative methods suggested and analyzed in this paper for solving extended general triequilibrium inclusions are the novel generalizations, improvements, refinements and modifications of Noor (three step) iterations [30–32], which include Ishikawa (two-step) iterations, Mann (one step)

iterations and Picard method as special cases. Using the technique and ideas of Ashish et. al. [2,3], Cho et al. [8], Cristescu et al. [10,11], Kwuni et al. [20], Mahato [22], Natrangan et al. [24], Noor et al. [47,48,53,54], Pamsang et al. [56], Rattanaseeha et al. [57], Suantai et al. [62], Tomar et al. [63], Trinh et al. [65] and Yadav et al. [67], one can explore the applications of these multi step methods for solving the general triequilibrium inclusions in the fixed point theory, fractal geometry, chaos theory, coding, number theory, spectral geometry, dynamical systems, complex analysis, nonlinear programming, graphics, artificial intelligence, control engineering, management sciences, stock exchange, regression and link prediction problems [60], financial mathematical [4], and computer aided design. Comparison of these new methods with other technique is an open problem, which needs further research efforts.

Acknowledgments. The authors sincerely thank their respected professors, teachers, students, colleagues, collaborators, referees, editors, managing editors and friends, who have contributed, directly or indirectly to this research. The authors are grateful to the referees for their valuable comments and suggestions for the improvements of the final version.

References

- [1] G.D. Anderson, M.K. Vamanamurthy and M. Vuorinen, Generalized convexity and inequalities, *J. Math. Anal. Appl.* 335 (2007), 1294-1308.
- [2] K. Ashish, M. Rani and R. Chugh, Julia sets and Mandelbrot sets in Noor orbit, *Appl. Math. Comput.* 228 (2014), 615-631.
- [3] K. Ashish, J. Cao and M.A. Noor, Stabilization of fixed points in chaotic maps using Noor orbit with applications in cardiac arrhythmia, *J. Appl. Anal. Comput.* 13 (2023), 2452-2470.
- [4] F. Al-Azemi and O. Calin, Asian options with harmonic average, *Appl. Math. Inform. Sci.* 9 (2015), 1-9.
- [5] M.I. Bloach, M.A. Noor and K.I. Noor, Iterative schemes for triequilibrium-like problems, *Int. J. Anal. Appl.* 19 (2021), 743-759.
- [6] M.I. Bloach, M.A. Noor and K.I. Noor, Well-posedness of triequilibrium-like problems, *Int. J. Anal. Appl.* 20 (2022), 1-10.

- [7] E. Blum and W.Oettli, From optimization and variational inequalities to equilibrium problems, *Math. Students* 63 (1994), 123–145.
- [8] S.Y. Cho, A.A. Shahid, W. Nazeer and S.M. Kang, Fixed point results for fractal generation in Noor orbit and s -convexity, *Springer Plus* 5 (2006), 18-43.
- [9] R.W. Cottle , J.-S. Pang and R.E. Stone, *The Linear Complementarity Problem*, SIAM Publ. Philadelphia, USA, 2009.
- [10] G. Cristescu and L. Lupsa, *Non-Connected Convexities and Applications*, Kluwer Academic Publishers, Dordrecht, Holland, 2002.
- [11] G. Cristescu and G. Mihail, Shape properties of Noor’s g -convex sets, *Proc. Twelfth Symposium of. Math. Appl.*, Timisoara, Romania, (2009), 1-13.
- [12] S. Dafermos, Sensitivity analysis in variational inequalities, *Math. Oper. Res.* 13 (1988), 421-434.
- [13] P. Dupuis and A. Nagurney, Dynamical systems and variational inequalities, *Anal. Oper. Res.* 44 (1993), 7-42.
- [14] R. Glowinski, J.L. Lions and R. Tremolieres, *Numerical Analysis of Variational Inequalities*, NorthHolland, Amsterdam, 1981.
- [15] N.K. Gupta, S. Varghese, M.A. Siddiqui and M.A. Noor, General mixed quasi equilibrium variational inequalities, *Global J. Pure Appl. Math.* 21 (2025), 407-418.
- [16] N.K. Gupta, M.A. Noor, K.I. Noor and S. Varghese, Iterative metthods for general bifunction equilibrium problems, *Global J. Pure Appl. Math.* 21 (2025), 991-1006.
- [17] S. Jabeen, J.E. Mac’ias-D’iaz, M.A. Noor, M.B. Khan and K.I. Noor, Design and convergence analysis of some implicit inertial methods for quasi-variational inequalities via the Wiener-Hopf equations, *Appl. Numer. Math.* 182 (2022), 76-86.
- [18] S. Jana and M.A. Noor, Mixed quasi-hemiequilibrium problems on Hadamard manifolds, *Int. J. Math. Stat. Oper. Res.* 5 (2025), 269-285
- [19] D. Kinderlehrer and G. Stampacchia, *An Introduction to Variational Inequalities and Their Applications*, SIAM, Philadelphia, USA, 2000.

- [20] Y.C. Kwuni, A.A. Shahid, W. Nazeer, S.I. Butt, M. Abbas and S.M. Kang, Tricorns and multicorns in Noor Orbit with s -convexity, *IEEE Access* 7 (2019), 2019.
- [21] J. Lions and G. Stampacchia, Variational inequalities, *Commun. Pure Appl. Math.* 20 (1967), 493-519.
- [22] N.K. Mahato, M.A. Noor and N.K. Sahu, Existence results for trifunction equilibrium problems and fixed point problems, *Anal. Math. Phys.* 9 (2019), 323–347.
- [23] A. Nagurney and D. Zhang, *Projected Dynamical Systems and Variational Inequalities with Applications*, Kluwer Academic Publishers, Boston, Dordrecht, London, 1996.
- [24] S.K. Natarajan and D. Negi, Green innovations utilizing fractal and power for solar panel optimization, in *Green Innovations for Industrial Development and Business Sustainability*. In: R.Sharma, G. Rana and S. Agarwal (Eds), CRC Press, Florida, Boca Raton, USA, 2024, pp.146-152.
- [25] C.P. Niculescu and L.E. Persson, *Convex Functions and Their Applications*, Springer-Verlag, New York, 2018.
- [26] M.A. Noor, *On Variational Inequalities*, PhD Thesis, Brunel University, London, U.K. 1975.
- [27] M.A. Noor, General variational inequalities, *Appl. Math. Lett.* 1 (1988), 119-121.
- [28] M.A. Noor, Quasi-variational inequalities, *Appl. Math. Lett.* 1 (1988), 367-370.
- [29] M.A. Noor, Sensitivity analysis for quasi-variational inequalities, *J. Optim. Theory Appl.* 95 (1997), 399-407.
- [30] M.A. Noor, New approximation schemes for general variational inequalities, *J. Math. Anal. Appl.* 251 (2000), 217-229.
- [31] M.A. Noor, Three-step iterative algorithms for multivalued quasi-variational inclusions, *J. Math. Anal. Appl.* 255 (2001), 589-604.
- [32] M.A. Noor, Some developments in general variational inequalities, *Appl. Math. Comput.* 152 (2004), 199-277.

- [33] M.A. Noor, Auxiliary principle technique for equilibrium problems, *J. Optim. Theory Appl.* 122 (2004), 371–386.
- [34] M.A. Noor, Multivalued general equilibrium problems, *J. Math. Anal. Appl.* 283 (2003), 140–149.
- [35] M.A. Noor, On a class of nonconvex equilibrium problems, *Appl. Math. Comput.* 157(2004), 553–566.
- [36] M.A. Noor, Generalized mixed quasi-equilibrium problems with trifunction, *Appl. Math. Lett.* 18 (2005), 695–700.
- [37] M.A. Noor and K.I. Noor, On equilibrium problems, *Appl. Math. E-Notes* 4 (2004), 125–132.
- [38] M.A. Noor, Fundamentals of equilibrium problems, *Math. Ineq. Appl.* 6 (2006), 529–566.
- [39] M.A. Noor, Differentiable nonconvex functions and general variational inequalities, *Appl. Math. Comput.* 199 (2008), 623–630.
- [40] M.A. Noor, Extended general variational inequalities, *Appl. Math. Lett.* 22 (2009), 182–185.
- [41] M.A. Noor and E. Al-Said, Change of variable method for generalized complementarity problems, *J. Optim. Theory Appl.* 100 (1999), 389–395.
- [42] M.A. Noor and K.I. Noor, Sensitivity analysis for quasi variational inclusions, *J. Math. Anal. Appl.* 236 (1999), 290–299.
- [43] M.A. Noor and K.I. Noor, Equilibrium problems and variational inequalities, *Mathematica (Cluj)*, 47 (2005), 89–100.
- [44] M.A. Noor and K.I. Noor, Dynamical system technique for solving quasi variational inequalities, *U.P.B. Sci. Bull., Ser. A* 84 (2022), 55–66.
- [45] M.A. Noor and K.I. Noor, New Inertial approximation schemes for general quasi variational inclusions, *Filomat* 36 (2022), 6071–6084.
- [46] M.A. Noor and K.I. Noor, General bivariational inclusions and iterative methods, *Int. J. Nonlinear Anal. Appl.* 14 (2023), 309–324.
- [47] M.A. Noor and K.I. Noor, Some new iterative schemes for solving general quasi variational inequalities, *Le Matematiche* 79 (2024), 327–370.

- [48] M.A. Noor and K.I. Noor, Some novel aspects and applications of Noor iterations and Noor orbits, *J. Adv. Math. Stud.* 17 (2024), 276-284.
- [49] M.A. Noor and K.I. Noor, General harmonic-like variational inequalities, *U.P.B. Sci. Bull., Ser. A* 87 (2025), 49-58.
- [50] M.A. Noor and K.I. Noor, Some new classes of general harmonic-like nonlinear equations, *General Math.* 33 (2025), 17-35.
- [51] M.A. Noor and K.I. Noor, Auxiliary principle technique for solving trifunction harmonic variational inequalities, *Rad Hazu. Matemat. Znanosti* 29 (2025), 261-281.
- [52] M.A. Noor and K.I. Noor, Harmonic-like bifunction equilibrium problems, *J. Adva. Math. Stud.* 19 (2026), in Press.
- [53] M.A. Noor, K.I. Noor and M.T. Rassias, New trends in general variational inequalities, *Acta Appl. Math.* 170 (2020), 981-1046.
- [54] M.A. Noor, K.I. Noor and M.T. Rassias, *General variational inequalities and optimization*. In: Geometry and Non-Convex Optimization (Panos M. Pardalos, Themistocles M. Rassias), Springer Optimization and Its Applications, Springer Nature, Switzerland, 223 (2025), 361-611.
- [55] M.A. Noor and W. Oettli, On general nonlinear complementarity problems and quasi equilibria, *Le Matematiche* 49 (1994), 313-331.
- [56] S. Paimsang, D. Yambangwai and T. Thainwan, A novel Noor iterative method of operators with property (E) as concerns convex programming applicable in signal recovery and polynomiography, *Math. Methods Appl. Sci.* 47 (2024), 9571-9588.
- [57] K. Rattanaseeha, S. Imnang, P. Inkrong and T. Thianwan, Novel Noor iterative methods for mixed type asymptotically nonexpansive mappings from the perspective of convex programming in hyperbolic spaces, *Int. J. Innov. Comput. Inform. Cont.* 19 (2023), 1717-1734.
- [58] S. Reich, Extension problems for accretive sets in Banach spaces, *J. Funct. Anal.* 26 (1977), 378—395.
- [59] S. Reich and S. Sabach, Three convergence theorems regarding iterative methods for solving equilibrium problems in reflexive Banach spaces, *Contemp. Math.* 568 (2012), 225-240.

- [60] P.H. Sach and Le Anh Tuan, Generalized equilibria with multivalued trifunctions and applications to uncertain vector equilibrium problems, *J. Optim. Theory Appl.* 208 (2026), 1-29.
- [61] G. Stampacchia, Formes bilineaires coercitives sur les ensembles convexes, *C.R. Acad. Sci. Paris* 258 (1964), 4413-4416.
- [62] S. Suantai, M.A. Noor, K. Kankam and P. Cholamjiak, Novel forward-backward algorithms for optimization and applications to compressive sensing and image inpainting, *Adva. Differ. Equations* 2021 (2021), 265.
- [63] A. Tomar, S. Antal, M. Sajid and D.J. Prajapati, Role of s-convexity in the generation of fractals as Julia and Mandelbrot sets via three-step fixed point iteration, *AIMS Math.* 10 (2025), 26077–26105.
- [64] D.V. Thong, X.-H. Li, S. Reich, Q.-L. Dong and D.H. Ngan, A new approach to the Korpelevich method for solving pseudomonotone equilibrium problems, *Numer. Alg.* 98 (2025), 719—741,
- [65] T.Q. Trinh and P.T. Vuong, The projection algorithm for inverse quasi-variational inequalities with applications to traffic assignment and network equilibrium control problems, *Optimization* 2024 (2024), 1-25.
- [66] Y.S. Xia and J. Wang, On the stability of globally projected dynamical systems, *J. Optim. Theory Appl.* 106 (2000), 129-150.
- [67] A. Yadav and K. Jha, Parrondo's paradox in the Noor logistic map, *Int. J. Adv. Res. Eng. Tech.* 7 (2016), 01-06.
- [68] Y. Zhang and G. Yu, Error bounds for inverse mixed quasi-variational inequality via generalized residual gap functions, *Asia-Pacific J. Oper. Res.* 39 (2022), 2150017.